Catalog Number: 100831, 100834, 191335, 195303

## Lysozyme

**CAS** #: 9001-63-2

Synonym: Muramidase

Physical Appearance: White to off white lyophilized crystalline powder

**Description:** Lysozyme (muramidase) hydrolyzes preferentially the b-1,4 glucosidic linkages between N-acetylmuramic acid and N-acetylglucosamine which occur in the mucopeptide cell wall structure of certain microorganisms, such as *Micrococcus lysodeikticus*.

In humans, Osserman, et al. <sup>14</sup> have indicated that lysozyme may be the mediator in the anti-tumor function of macrophages which, it has been shown, secrete the enzyme. <sup>8</sup> There is evidence that cartilage lysozyme has a role in cartilage calcification. <sup>12</sup>

The enzyme has been used for lysing *E. coli* and *Streptomycetes* for extraction purposes<sup>9</sup> such as extracting group specific antigen. <sup>18</sup> It would appear that lysozyme may act as a germinative agent of bacterial spores. <sup>2,7</sup>

Molecular weight: 14,388 (Chicken<sup>11</sup>); 17,000 (human).

**Optimum pH:** 9.2.<sup>5</sup>

**Extinction coefficient:**  $= 26.4.^3$ 

**Isoelectric point:** pH 11.0.<sup>1</sup>

**Inhibitors:** The enzyme is inhibited by surface-active reagents such as dodecyl sulfate, alcohols and fatty acids. <sup>17</sup> Imidazole and indole derivatives are inhibitors via formation of change-transfer complexes. <sup>15</sup>

**Stability:** Solutions at pH 4-5 are stable for several weeks refrigerated and for days at ambient temperatures.

**Assay:** Mintz, et al. <sup>13</sup> describe a sensitive fluorimetric assay.

## **Unit Definitions:**

*From Chicken:* One unit will cause a decrease in A450 of 0.001 per minute at pH 6.24 and 25°C using Micrococcus lysodeikticus as substrate.

Shugar Unit: <sup>16</sup> One unit is defined as the amount of enzyme that will digest powdered cells of Micrococcus lysodeikticus, causing a decrease in absorbancy of 0.001 per minute at 37°C, pH 7.0

## **Availability:**

Catalog Number	Description	Size
100831		100 mg
	crystallized; Activity: 20,000 - 25,000 units/mg	500 mg

	protein	1 g 5 g 10 g 25 g 100 g
100834	Lysozyme, from chicken egg white, 2X crystallized; Activity: > 9,000 units/mg protein	1 g 5 g 10 g
195303	Lysozyme, from chicken egg white, Type VI, 3X crystallized; Activity: ~60,000 units/mg protein	1 g 5 g 25 g
191335	Lysozyme, from human neutrophils; Activity: ~30,000 Shugar units/mg protein	100 ug

## **References:**

- 1. Alderton, G., Ward, W., and Febold, H., "Isolation of Lysozyme from Egg White," *J. Biol. Chem.*, v. 157, 43 (1945).
- 2. Ando, Y., "The Effect of Lysozyme on Ionic Forms of Spores of *Clostridium perfringens* Type A," *J. Bacteriol.*, **v. 122**, 794 (1975).
- 3. Aune, K., and Tanford, C., "Thermodynamics of the Denaturation of Lysozyme by Guanidine Hydrochloride. I. Dependence on pH at 25°C.," *Biochem.*, v. 8, 4579 (1969).
- 4. Dahlquist, F., Borders, C., Jacobson, G., and Raftery, M., "The Stereospecificity of Human, Hen and Papaya Lysozymes," *Biochem.*, v. 8, 694 (1969).
- 5. Davies, A., Neuberger, A., and Wilson, B., "The Dependence of Lysozyme Activity on pH and Ionic Strength," *Biochim. Biophys. Acta*, v. 178, 294 (1969).
- 6. Dubois, T., Guillard, R., Prieels, J., and Perraudin, J., "Comparison between the Folding of Reduced Hen Egg White Lysozyme and that of Reduced Human Milk Lysozyme," *Biochem.*, v. 21, 6516 (1982).
- 7. Duncan, C., Labbe, R., and Reich, R., "Germination of Heat- and Alkali-Altered Spores of *Clostridium perfringens* Type A by Lysozyme and and Initiation Protein,, " *J. Bacteriol.*, **v. 109**, 550 (1972).
- 8. Gordon, S., Todd, J., and Cohn, Z., "In Vitro Synthesis and Secretion of Lysozyme by Mononuclear Phagocytes," *J. Exp. Med.*, v. 139, 1228 (1974).
- 9. Haas, M., and Dowding, J., "Aminoglycoside-Modifying Enzymes," in *Methods in Enzymology*, **XLIII**, (Hash, J., ed.), Academic Press, NY, 621 (1965).
- 10. Hankiewicz, J., and Swierczek, E., "Lysozyme in Human Body Fluids," Clin. Chim. Acta, v. 57, 205 (1974).
- 11. Jollés, P., "Lysozyme," in *The Enzymes*, **IV**, 2nd Ed., (Boyer, P., Lardy, H., and Myrback, K., eds.), Academic Press, NY, 431 (1960).
- 12. Kuettner, K., Sorgente, N., Croxen, R., Howell, D., and Pita, J., "Lysozyme in Preosseous Cartilage. VII. Evidence for Physiological Role of Lysozyme in Normal Endochondral Calcificaion," *Biochim. Biophys. Acta*, v. 372, 335 (1974).
- 13. Mintz, G., Herbold, D., and Glaser, L., "A Fluorescent Assay for Bacterial Cell Wall Lytic Enzymes," *Anal. Biochem.*, v. 66, 272 (1975).
- 14. Osserman, E., Kockars, M., Halper, J., and Fischel, R., "Effects of Lysozyme on Normal and Transformed Cells," *Nature*, v. 243, 331 (1973).
- 15. Shinitzky, M., Katchalski, E., Grisaro, V., and Sharon, N., "Inhibition of Lysozyme by Imidazole and Indole Derivatives," *Arch. Biochem. Biophys.*, v. 116, 332 (1966).
- 16. Shugar, D., "Measurement of Lysozyme Activity and the Ultra Violet Inactivation of Lysozyme," *Biochim. Biophys. Acta*, **v. 8**, 302 (1952).
- 17. Smith, G., and Stoker, C., "Inhibition of Crystalline Lysozyme," Arch. Biochem., v. 21, 383 (1949).
- 18. Watson, B., Moellering, R., and Kunz, L., "Identification of *Streptococci*: Use of Lysozyme and *Streptomyces albus* Filtrate in the Preparation of Extracts for Lancefield Grouping," *J. Clin. Microbiol.*, v. 1, 274 (1975).

