Nuclear GreenTM DCS1

Ordering Information	Storage Conditions
Product Number: 17550 (5 mM)	Keep at -20 °C and avoid exposure to light

Spectral Properties

Ex/Em = 515/530 nm




Figure 1. Excitation and emission spectra for the Nuclear Green™ DCS1 bound to DNA in PBS (pH 7.4).

Biological Applications

Our Nuclear GreenTM DCS1 is a non-fluorescent, DNA-selective and cell impermeant green fluorescent dye for analyzing DNA content in dead cells. The Nuclear GreenTM DCS1 has its fluorescence significantly enhanced upon binding to double-stranded DNA. It can be used in fluorescence imaging, microplate and flow cytometry applications. This DNA-binding dye might be used for multicolor analysis of dead cells.

Sample Protocol for Staining Cells

Caution: The following protocol can be adapted for most cell types. Growth medium, cell density, the presence of other cell types and factors may influence staining. Residual detergent on glassware may also affect staining of many organisms, and cause brightly stained material to appear in solutions with or without cells present.

- 1. Add Nuclear GreenTM DCS1 (2 to 10 μ M) into the fixed, dead or apoptotic cells (either suspension or adherent) and stain the cells for 15 to 60 minutes. In initial experiments, it may be best to try several dye concentrations to determine the optimal concentration that yields the desired result. High dye concentration tends to cause nonspecific staining of other cellular structures.
- 2. Directly analyze the cellular staining with fluorescence microscopy, fluorescence microplate reader, or flow cytometry.

Disclaimer: This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact our technical service representative for more information.