



15 µg	Lot: 0041202	Exp: 2/14
100 µg/ml	Store at -20°C	

Description: NIH 3T3 (mouse embryonic fibroblast cell line) genomic DNA.

Source: NIH 3T3 (mouse embryonic fibroblast) cells are grown to confluency in DMEM plus 10% fetal bovine serum. Genomic DNA was isolated by a standard genomic purification protocol (1), phenol extracted and equilibrated to 10 mM Tris-HCI (pH 7.5) and 1 mM EDTA.



BioLabs

1-800-632-7799

info@neb.com

www.neb.com

BioLabs.

1-800-632-7799

info@neb.com

www.neb.com

- A control for Methylation-Specific PCR (MSP) (2), Bisulfite sequencing, Methylationsensitive Single-Nucleotide Primer Extension (Ms- SNuPE), Combined Bisulfite Restriciton Analysis (COBRA), Bisulfite treatment and PCR-Single-Strand Conformation Polymorphism Analysis (Bisulfite-PCR-SSCP/ BiPS).
- PCR, SNP Analysis, Southern Blotting
- Genomic DNA library construction

Supplied in: 10 mM Tris-HCl (pH 7.5) and 1 mM EDTA. Avoid multiple freeze/thaw cycles.

Quality Assurance: Purified free of contaminating proteins and RNA.

A260/280 Ratio: 1.90

Reference:

- Sambrook, J. and Russell, D. (2001) *Molecular Cloning: A Laboratory Manual,* (3rd ed.), (pp. 6.4–6.12). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
- Herman, J.G. and Baylin, S.B. (1996). U.S. Patent No. 5,786,146. John Hopkins University School of Medicine.

CERTIFICATE OF ANALYSIS



N4004S 004120214021

N4004S

15 µg	Lot: 0041202	Exp: 2/14
100 µg/ml	Store at -20°C	

Description: NIH 3T3 (mouse embryonic fibroblast cell line) genomic DNA.

Source: NIH 3T3 (mouse embryonic fibroblast) cells are grown to confluency in DMEM plus 10% fetal bovine serum. Genomic DNA was isolated by a standard genomic purification protocol (1), phenol extracted and equilibrated to 10 mM Tris-HCI (pH 7.5) and 1 mM EDTA.

Applications:

- A control for Methylation-Specific PCR (MSP) (2), Bisulfite sequencing, Methylationsensitive Single-Nucleotide Primer Extension (Ms- SNuPE), Combined Bisulfite Restriciton Analysis (COBRA), Bisulfite treatment and PCR-Single-Strand Conformation Polymorphism Analysis (Bisulfite-PCR-SSCP/ BiPS).
- PCR, SNP Analysis, Southern Blotting
- Genomic DNA library construction

Supplied in: 10 mM Tris-HCl (pH 7.5) and 1 mM EDTA. Avoid multiple freeze/thaw cycles.

Quality Assurance: Purified free of contaminating proteins and RNA.

A260/280 Ratio: 1.90

Reference:

- Sambrook, J. and Russell, D. (2001) *Molecular Cloning: A Laboratory Manual,* (3rd ed.), (pp. 6.4–6.12). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
- Herman, J.G. and Baylin, S.B. (1996). U.S. Patent No. 5,786,146. John Hopkins University School of Medicine.