

# HincII



1-800-632-7799  
info@neb.com  
www.neb.com



R0103S 046120814081

## R0103S



1,000 units 10,000 U/ml Lot: 0461208

RECOMBINANT Store at -20°C Exp: 8/14

### Recognition Site:

5'... G T Y R A C ... 3'  
3'... C A R Y T G ... 5'

**Single Letter Code:** R = A or G, Y = C or T

**Source:** An *E. coli* strain that carries the cloned HincII gene from *Haemophilus influenzae* Rc (ATCC 49699)

Supplied in: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

**Reagents Supplied with Enzyme:**  
10X NEBuffer 3, 100X BSA.

**Reaction Conditions:** 1X NEBuffer 3, supplemented with 100 µg/ml BSA. Incubate at 37°C.

### 1X NEBuffer 3:

100 mM NaCl  
50 mM Tris-HCl  
10 mM MgCl<sub>2</sub>  
1 mM dithiothreitol  
pH 7.9 @ 25°C

**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

### Diluent Compatibility:

Diluent Buffer B  
300 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM dithiothreitol, 500 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C)

### Quality Control Assays

**Ligation:** After 10-fold overdigestion with HincII, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of DNA and 100 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

**Exonuclease Activity:** Incubation of 100 units of enzyme with 1 µg sonicated <sup>3</sup>H DNA (10<sup>5</sup> cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

### Enzyme Properties

#### Activity in NEBuffers:

NEBuffer 1 75%  
NEBuffer 2 100%  
NEBuffer 3 **100%**  
NEBuffer 4 100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

**Survival in a Reaction:** A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

**Heat Inactivation:** 65°C for 20 minutes.

**Notes:** A 100-fold overdigestion of HincII in NEBuffer 4 produces star activity.

Cleavage of mammalian genomic DNA is blocked by some combinations of overlapping CpG methylation.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS

# HincII



1-800-632-7799  
info@neb.com  
www.neb.com



R0103S 046120814081

## R0103S



1,000 units 10,000 U/ml Lot: 0461208

RECOMBINANT Store at -20°C Exp: 8/14

### Recognition Site:

5'... G T Y R A C ... 3'  
3'... C A R Y T G ... 5'

**Single Letter Code:** R = A or G, Y = C or T

**Source:** An *E. coli* strain that carries the cloned HincII gene from *Haemophilus influenzae* Rc (ATCC 49699)

Supplied in: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

**Reagents Supplied with Enzyme:**  
10X NEBuffer 3, 100X BSA.

**Reaction Conditions:** 1X NEBuffer 3, supplemented with 100 µg/ml BSA. Incubate at 37°C.

### 1X NEBuffer 3:

100 mM NaCl  
50 mM Tris-HCl  
10 mM MgCl<sub>2</sub>  
1 mM dithiothreitol  
pH 7.9 @ 25°C

**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

### Diluent Compatibility:

Diluent Buffer B  
300 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM dithiothreitol, 500 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C)

### Quality Control Assays

**Ligation:** After 10-fold overdigestion with HincII, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of DNA and 100 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

**Exonuclease Activity:** Incubation of 100 units of enzyme with 1 µg sonicated <sup>3</sup>H DNA (10<sup>5</sup> cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

### Enzyme Properties

#### Activity in NEBuffers:

NEBuffer 1 75%  
NEBuffer 2 100%  
NEBuffer 3 **100%**  
NEBuffer 4 100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

**Survival in a Reaction:** A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

**Heat Inactivation:** 65°C for 20 minutes.

**Notes:** A 100-fold overdigestion of HincII in NEBuffer 4 produces star activity.

Cleavage of mammalian genomic DNA is blocked by some combinations of overlapping CpG methylation.

= Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS