



EarI









R0528S 045120914091

R0528S



1-800-632-7799
info@neb.com
www.neb.com

500 units
20,000 U/ml
Lot: 0451209

RECOMBINANT
Store at -20°C
Exp: 9/14

Recognition Site:

5'...CTCTTC(N)₁▼...3'
3'...GAGAAG(N)₄▲...5'

Source: An *E. coli* strain that carries the cloned Earl gene from *Enterobacter aerogenes* (C. Polisson)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

New Reaction Buffer

EarI



R0528S 045120914091

R0528S



1-800-632-7799
info@neb.com
www.neb.com









500 units
20,000 U/ml
Lot: 0451209

RECOMBINANT
Store at -20°C
Exp: 9/14

Recognition Site:

5'...CTCTTC(N)₁▼...3'
3'...GAGAAG(N)₄▲...5'

Source: An *E. coli* strain that carries the cloned Earl gene from *Enterobacter aerogenes* (C. Polisson)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

New Reaction Buffer

Reagents Supplied with Enzyme:
10X NEBuffer 4.

Reaction Conditions: 1X NEBuffer 4.
Incubate at 37°C.

1X NEBuffer 4:
50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer A
50 mM KCl, 10 mM Tris-HCl, 0.1 mM EDTA,
1 mM DTT, 200 µg/ml BSA and 50% glycerol
(pH 7.4 @ 25°C).

Quality Control Assays

Ligation: After 10-fold overdigestion with Earl, approximately 75% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, approximately 75% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA and 50 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 40 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Enzyme Properties

Activity in NEBuffers:


NEBuffer 1	100%
NEBuffer 2	100%
NEBuffer 3	50%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 20 units of enzyme were inactivated by incubation at 65°C for 20 minutes.

Note: Cleavage of mammalian genomic DNA is impaired by overlapping CpG methylation.

 = Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS

Quality Control Assays

Ligation: After 10-fold overdigestion with Earl, approximately 75% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, approximately 75% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA and 50 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 40 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.1% radioactivity.

Enzyme Properties

Activity in NEBuffers:


NEBuffer 1	100%
NEBuffer 2	100%
NEBuffer 3	50%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 20 units of enzyme were inactivated by incubation at 65°C for 20 minutes.

Note: Cleavage of mammalian genomic DNA is impaired by overlapping CpG methylation.

 = Time-Saver™ Qualified (See www.neb.com for details).

CERTIFICATE OF ANALYSIS