

BsaI



1-800-632-7799
info@neb.com
www.neb.com



R0535S 034120714071

R0535S



1,000 units 10,000 U/ml Lot: 0341207

RECOMBINANT Store at -20°C Exp: 7/14

Recognition Site:

5'...GGTCTC(N)₁...3'
3'...CCAGAG(N)₅...5'

Source: An *E. coli* strain that carries the cloned BsaI gene from *Bacillus stearothermophilus* 6-55 (Z. Chen)

Reaction Temperature now 37°C

BsaI



1-800-632-7799
info@neb.com
www.neb.com



R0535S 034120714071

R0535S



1,000 units 10,000 U/ml Lot: 0341207

RECOMBINANT Store at -20°C Exp: 7/14

Recognition Site:

5'...GGTCTC(N)₁...3'
3'...CCAGAG(N)₅...5'

Source: An *E. coli* strain that carries the cloned BsaI gene from *Bacillus stearothermophilus* 6-55 (Z. Chen)

Reaction Temperature now 37°C

Supplied in: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X NEBuffer 4, 100X BSA.

Reaction Conditions:

1X NEBuffer 4. Supplement with 100 µg/ml BSA. Incubate at 37°C.

1X NEBuffer 4:

50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility:

Diluent Buffer B
300 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA,
1 mM DTT, 500 µg/ml BSA and 50% glycerol
(pH 7.4 @ 25°C)

Supplied in: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X NEBuffer 4, 100X BSA.

Reaction Conditions:

1X NEBuffer 4. Supplement with 100 µg/ml BSA. Incubate at 37°C.

1X NEBuffer 4:

50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility:

Diluent Buffer B
300 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA,
1 mM DTT, 500 µg/ml BSA and 50% glycerol
(pH 7.4 @ 25°C)

Quality Control Assays

Ligation: After 10-fold overdigestion with BsaI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 200 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 100 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.05% radioactivity.

Endonuclease Activity: Incubation of 50 units of enzyme with 1 µg φX174 RF I DNA for 4 hours 37°C in 50 µl reaction buffer resulted in < 5% conversion to RF II.

Quality Control Assays

Ligation: After 10-fold overdigestion with BsaI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1–2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 200 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

Exonuclease Activity: Incubation of 100 units of enzyme with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.05% radioactivity.

Endonuclease Activity: Incubation of 50 units of enzyme with 1 µg φX174 RF I DNA for 4 hours 37°C in 50 µl reaction buffer resulted in < 5% conversion to RF II.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1	75%
NEBuffer 2	75%
NEBuffer 3	100%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: Not recommended for digest over 1 hour.

Heat Inactivation: 65°C for 20 minutes.

Plasmid Cleavage: Number of units required to cleave 1 µg of supercoiled plasmid DNA in one hour: 2 units.

Note: Blocked by overlapping *dcm* methylation. Cleavage of mammalian genomic DNA is blocked by some combinations of overlapping CpG methylation.

The addition of BSA to the restriction digest allows BsaI to be used at 37°C. This is also true for older lots of BsaI as there has been no formulation change in the enzyme. Activity at 50°C is 100%.

CERTIFICATE OF ANALYSIS

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1	75%
NEBuffer 2	75%
NEBuffer 3	100%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: Not recommended for digest over 1 hour.

Heat Inactivation: 65°C for 20 minutes.

Plasmid Cleavage: Number of units required to cleave 1 µg of supercoiled plasmid DNA in one hour: 2 units.

Note: Blocked by overlapping *dcm* methylation. Cleavage of mammalian genomic DNA is blocked by some combinations of overlapping CpG methylation.

The addition of BSA to the restriction digest allows BsaI to be used at 37°C. This is also true for older lots of BsaI as there has been no formulation change in the enzyme. Activity at 50°C is 100%.

CERTIFICATE OF ANALYSIS