

PspOMI



1-800-632-7799
info@neb.com
www.neb.com



R0653S 006120914091

R0653S



1,500 units **20,000 U/ml** **Lot: 0061209**
RECOMBINANT **Store at -20°C** **Exp: 9/14**

Recognition Site:

5'...GGGCC...3'
3'...CCCGG...5'

Source: An *E. coli* strain that carries the cloned PspOMI gene from *Pseudomonas* species OM2164

New Storage Conditions

PspOMI



1-800-632-7799
info@neb.com
www.neb.com



R0653S 006120914091

R0653S



1,500 units **20,000 U/ml** **Lot: 0061209**
RECOMBINANT **Store at -20°C** **Exp: 9/14**

Recognition Site:

5'...GGGCC...3'
3'...CCCGG...5'

Source: An *E. coli* strain that carries the cloned PspOMI gene from *Pseudomonas* species OM2164

New Storage Conditions

Supplied in: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 500 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 4

Reaction Conditions: 1X NEBuffer 4
Incubate at 37°C.

1X NEBuffer 4:
20 mM Tris-acetate
10 mM magnesium acetate
50 mM potassium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer B
300 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM dithiothreitol, 500 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C)

Supplied in: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 500 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer 4

Reaction Conditions: 1X NEBuffer 4
Incubate at 37°C.

1X NEBuffer 4:
20 mM Tris-acetate
10 mM magnesium acetate
50 mM potassium acetate
1 mM DTT
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Diluent Compatibility: Diluent Buffer B
300 mM NaCl, 10 mM Tris-HCl, 0.1 mM EDTA, 1 mM dithiothreitol, 500 µg/ml BSA and 50% glycerol (pH 7.4 @ 25°C)

Quality Control Assays

Ligation: After 20-fold overdigestion with PspOMI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1-2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 200 units of enzyme incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by gel electrophoresis.

Exonuclease Activity: Incubation of a 50 µl reaction containing 200 units of PspOMI with 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA (200,000 cpm/µg) for 4 hours at 37°C released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 µl reaction containing 200 units of PspOMI with 1 µg of ϕX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

Quality Control Assays

Ligation: After 20-fold overdigestion with PspOMI, > 95% of the DNA fragments can be ligated with T4 DNA Ligase (at a 5' termini concentration of 1-2 µM) at 16°C. Of these ligated fragments, > 95% can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 200 units of enzyme incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by gel electrophoresis.

Exonuclease Activity: Incubation of a 50 µl reaction containing 200 units of PspOMI with 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA (200,000 cpm/µg) for 4 hours at 37°C released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 µl reaction containing 200 units of PspOMI with 1 µg of ϕX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

Enzyme Properties

Activity in NEBuffers:
NEBuffer 1 25%
NEBuffer 2 25%
NEBuffer 3 10%
NEBuffer 4 **100%**

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 65°C for 20 minutes.

Notes: PspOMI is an isoschizomer of Bsp120I.

Impaired by some combinations of overlapping *dcm* methylation. Cleavage of mammalian genomic DNA is blocked by overlapping CpG methylation.

Companion Products:

dam⁻/*dcm*⁻ Competent *E. coli*
#C2925H 20 transformation reactions
#C2925I 24 transformation reactions

CERTIFICATE OF ANALYSIS

Enzyme Properties

Activity in NEBuffers:
NEBuffer 1 25%
NEBuffer 2 25%
NEBuffer 3 10%
NEBuffer 4 **100%**

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.13 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 65°C for 20 minutes.

Notes: PspOMI is an isoschizomer of Bsp120I.

Impaired by some combinations of overlapping *dcm* methylation. Cleavage of mammalian genomic DNA is blocked by overlapping CpG methylation.

Companion Products:

dam⁻/*dcm*⁻ Competent *E. coli*
#C2925H 20 transformation reactions
#C2925I 24 transformation reactions

CERTIFICATE OF ANALYSIS