


# **Anti-Human CD19 FITC**

Catalog Number: 11-0199 Also Known As:Leu-12 RUO: For Research Use Only. Not for use in diagnostic procedures.



### **Product Information**

Contents: Anti-Human CD19 FITC

REF Catalog Number: 11-0199 Clone: HIB19 Concentration: 5 uL (1 ug)/test Host/Isotype: Mouse IgG1, kappa HLDA Workshop: V CD19.11 **Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

**Temperature Limitation:** Store at 2-8°C. Do not freeze.

Light sensitive material.

 Image: Batch Code: Refer to Vial

Use By: Refer to Vial

## Description

The HIB19 monoclonal antibody reacts with human CD19, a 95 kDa transmembrane glycoprotein. CD19 is expressed by B cells during all stages of development excluding the terminally differentiated plasma cells. Follicular dendritic cells also express CD19. Together CD21, CD81, Leu13, MHC class II, and CD19 form a multimolecular complex that associates with BCR. Signaling through CD19 induces tyrosine phosphorylation, calcium flux and proliferation of B cells.

## **Applications Reported**

The HIB19 antibody has been reported for use in flow cytometric analysis.

## **Applications Tested**

This HIB19 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5  $\mu$ L (1  $\mu$ g) per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test.

## References

Knapp, W., B. Dorken, et al. eds. (1989). Leucocyte Typing IV: White Cell Differentiation Antigens. Oxford University Press. New York.

Schlossman, S., L. Bloumsell, et al. eds (1995). Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press. New York.

## **Related Products**

11-4714 Mouse IgG1 K Isotype Control FITC (P3.6.2.1) 12-0038 Anti-Human CD3 PE (UCHT1)