


Anti-Human CD42a FITC

Catalog Number: 11-0428

Also known as: Glycoprotein IX, GP9

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human platelets with Anti-Human CD41 PE (cat. 12-0419) and Mouse IgG1 K Isotype Control FITC (cat. 11-4714) (left) or Anti-Human CD42a FITC (right). Total viable cells were used for analysis.

Product Information



Contents: Anti-Human CD42a FITC

Catalog Number: 11-0428

Clone: GR-P

Concentration: 5 μ L (0.25 μ g)/test

Host/Isotype: Mouse IgG1, k



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material.



Batch Code: Refer to vial



Use By: Refer to vial

Description

This GR-P monoclonal antibody reacts with human CD42a, which is also known as glycoprotein IX (GPIX). CD42a is a 22-kDa transmembrane protein that is constitutively expressed on platelets and megakaryocytes. CD42a associates with CD42b, CD42c, and CD42d to form the GPIb-V-IX complex, which binds the von Willebrand factor, an endothelial adhesion protein, to mediate platelet adhesion and activation. This binding results in platelet aggregation at sites of vascular injury. This receptor complex has also been demonstrated to bind thrombin, factors XI and XII, P-selectin, and Mac-1. Finally, a point mutation within the leucine-rich motif of CD42a has been linked to Bernard-Soulier syndrome.

Applications Reported

This GR-P antibody has been reported for use in flow cytometric analysis.

Applications Tested

This GR-P antibody has been pre-titrated and tested by flow cytometric analysis of human platelets. This can be used at 5 μ L (0.25 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Eisbacher M, Khachigian LM, Khin TH, Holmes ML, Chong BH. Inducible expression of the megakaryocyte-specific gene glycoprotein IX is mediated through an Ets binding site and involves upstream activation of extracellular signal-regulated kinase. *Cell Growth Differ.* 2001 Aug;12(8):435-45. (**GR-P**, FC)

Canobbio I, Balduini C, Torti M. Signalling through the platelet glycoprotein Ib-V-IX complex. *Cell Signal.* 2004 Dec;16(12):1329-44.

Slupsky JR, Kamiguti AS, Rhodes NP, Cawley JC, Shaw AR, Zuzel M. The platelet antigens CD9, CD42 and integrin alpha IIb beta IIIa can be topographically associated and transduce functionally similar signals. *Eur J Biochem.* 1997 Feb 15;244(1):168-75.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com

Anti-Human CD42a FITC

Catalog Number: 11-0428

Also known as: Glycoprotein IX, GP9

RUO: For Research Use Only. Not for use in diagnostic procedures.

Hickey MJ, Williams SA, Roth GJ. Human platelet glycoprotein IX: an adhesive prototype of leucine-rich glycoproteins with flank-center-flank structures. Proc Natl Acad Sci U S A. 1989 Sep;86(17):6773-7.

Related Products

11-4714 Mouse IgG1 K Isotype Control FITC (P3.6.2.8.1)

12-0419 Anti-Human CD41a PE (HIP8)

12-0429 Anti-Human CD42b PE (HIP1)