


Anti-Mouse CD107b (LAMP-2) FITC

Catalog Number: 11-1072

Also Known As: LAMP2, lysosomal-associated membrane protein 2

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of thioglycolate-induced peritoneal exudate cells with 0.25 ug of Rat IgG2a K Isotype Control FITC (cat. 11-4321) (open histogram) or 0.25 ug of Anti-Mouse CD107b (LAMP-2) FITC (filled histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD107b (LAMP-2) FITC

REF Catalog Number: 11-1072

Clone: eBioABL-93 (ABL-93)

Concentration: 0.5 mg/mL

Host/Isotype: Rat IgG2a, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.



Batch Code: Refer to Vial



Use By: Refer to Vial



Contains sodium azide

Description

The eBioABL-93 monoclonal antibody reacts with mouse CD107b, also known as lysosomal-associated membrane protein-2 (LAMP-2). CD107b is a highly glycosylated protein of approximately 120 kDa that is expressed intracellularly in lysosomal membranes. It is also transiently expressed on the surface of cytolytic T cells during degranulation, but to a lesser extent than CD107a. It has been reported that CD107b may be identical to the mouse Mac-3 antigen.

Applications Reported

This eBioABL-93 (ABL-93) antibody has been reported for use in intracellular staining followed by flow cytometric analysis. It is also reported for use in surface staining in a flow cytometric based degranulation assay.

Applications Tested

This eBioABL-93 (ABL-93) antibody has been tested by intracellular staining and flow cytometric analysis of thioglycolate-induced peritoneal-exudate cells. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Betts MR, Koup RA. Detection of T-cell degranulation: CD107a and b. *Methods Cell Biol.* 2004;75:497-512.

Tritel M, Stoddard AM, Flynn BJ, Darrah PA, Wu CY, Wille U, Shah JA, Huang Y, Xu L, Betts MR, Nabel GJ, Seder RA. Prime-boost vaccination with HIV-1 Gag protein and cytosine phosphate guanosine oligodeoxynucleotide, followed by adenovirus, induces sustained and robust humoral and cellular immune responses. *J Immunol.* 2003 Sep 1;171(5):2538-47. (ABL-93, degranulation assay, PubMed)

Cha Y, Holland SM, August JT. The cDNA sequence of mouse LAMP-2. Evidence for two classes of lysosomal membrane glycoproteins. *J Biol Chem.* 1990 Mar 25;265(9):5008-13. (ABL-93, affinity-chromatography, PubMed)

Chen JW, Chen GL, D'Souza MP, Murphy TL, August JT. Lysosomal membrane glycoproteins: properties of LAMP-1 and LAMP-2. *Biochem Soc Symp.* 1986;51:97-112.

Chen JW, Murphy TL, Willingham MC, Pastan I, August JT. Identification of two lysosomal membrane glycoproteins. *J Cell Biol.* 1985 Jul;101(1):85-95. (ABL-93, IP, IHC, PubMed)

Related Products

11-1078 Anti-Human CD107b (LAMP-2) FITC (eBioH4B4 (H4B4))
11-4321 Rat IgG2a K Isotype Control FITC (eBR2a)
53-1071 Anti-Mouse CD107a (LAMP-1) Alexa Fluor® 488 (eBio1D4B (1D4B))
88-8823 Fixation & Permeabilization Buffers

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com