


Anti-Human CD116 FITC

Catalog Number: 11-1169

Also Known As: GM-CSF Receptor alpha, CSF2RA

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Mouse IgG1 kappa Isotype Control FITC (cat. 11-4714) (open histogram) or Anti-Human CD116 FITC (filled histogram). Cells in the monocyte gate were used for analysis.

Product Information

Contents: Anti-Human CD116 FITC

REF **Catalog Number:** 11-1169

Clone: 4H1

Concentration: 5 µL (1 µg)/test

Host/Isotype: Mouse IgG1, kappa

HLDA Workshop: N/A

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

Description

The 4H1 monoclonal antibody reacts with the human CD116 molecule, the α subunit of GM-CSF receptor. The α subunit associates with the common β chain (CD131) to form the high affinity receptor for GM-CSF. The GM-CSFR α chain is expressed by granulocytes, monocytes, endothelial cells, fibroblasts and some tumor cells.

Applications Reported

4H1 has been reported for use in flow cytometric analysis.

Applications Tested

This 4H1 antibody has been pre-titrated and tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at 5 µL (1 µg) per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Sun, Q., K. Jones, et al. (1999). Simultaneous antagonism of interleukin-5, granulocyte-macrophage colony-stimulating factor, and interleukin-3 stimulation of human eosinophils by targeting the common cytokine binding site of their receptors. *Blood* 94(6): 1943-51.

Woodcock, J. M., B. J. McClure, et al. (1997). The human granulocyte-macrophage colony-stimulating factor (GM-CSF) receptor exists as a preformed receptor complex that can be activated by GM-CSF, interleukin-3, or interleukin-5. *Blood* 90(8): 3005-17.

Lopez, A. F., M. A. Vadas, et al. (1991). Interleukin-5, interleukin-3, and granulocyte-macrophage colony-stimulating factor cross-compete for binding to cell surface receptors on human eosinophils. *J Biol Chem* 266(36): 24741-7.

Related Products

11-4714 Mouse IgG1 K Isotype Control FITC (P3.6.2.1)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com