


Anti-Mouse NK1.1 FITC

Catalog Number: 11-5941

Also known as: CD161, NKR-P1C, Ly-55

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57Bl/6 splenocytes with Anti-Mouse CD49b (Integrin alpha 2) eFluor[®] 450 (cat. 48-5971) and 0.25 ug of Mouse IgG2a K Isotype Control FITC (cat. 11-4724) (left) or 0.25 ug of Anti-Mouse NK1.1 FITC (right). Cells in the lymphocyte gate were used for analysis.

Product Information



Contents: Anti-Mouse NK1.1 FITC

Catalog Number: 11-5941

Clone: PK136

Concentration: 0.5 mg/mL

Host/Isotype: Mouse IgG2a, kappa



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Caution, contains Azide



Description

The PK136 monoclonal antibody reacts with mouse NK1.1, an antigen expressed by natural killer cells and a subset of T cells in the NK1.1 mouse strains including C57BL and NZB. Several commonly used laboratory mouse strains such as BALB/c, SJL, AKR, CBA, C3H and A do not express the NK1.1 antigen. For detection of NK cells in these strains the monoclonal antibody DX5 14-5971 should be used. Simultaneous staining of C57BL/6 spleen cells with PK136 and DX5 reveals coexpression of both markers by a majority of cells as well as presence of small populations of DX5+PK136- and DX5-PK136+ cells.

Applications Reported

PK136 has been reported for use in flow cytometric analysis.

Applications Tested

The PK136 antibody has been tested by flow cytometric analysis of C57BL/6 mouse splenocyte suspensions and can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Kitaichi N, Kotake S, Morohashi T, Onoe K, Ohno S, Taylor AW. Diminution of experimental autoimmune uveoretinitis (EAU) in mice depleted of NK cells. *J Leukoc Biol.* 2002 Dec;72(6):1117-21. (**PK136**, in vivo depletion, PubMed)

Koo, G. C. and J. R. Peppard. Establishment of monoclonal anti-Nk-1.1 antibody. *Hybridoma* 1984. 3(3): 301-3.

Related Products

11-4724 Mouse IgG2a K Isotype Control FITC

12-0031 Anti-Mouse CD3e PE (145-2C11)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Mouse NK1.1 FITC

Catalog Number: 11-5941

Also known as: CD161, NKR-P1C, Ly-55

RUO: For Research Use Only. Not for use in diagnostic procedures.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com