

Anti-Human CD38 Purified

Catalog Number: 14-0389

Also Known As: ADP-Ribosyl Cyclase, Cyclic ADP-Ribose Hydrolase

RUO: For Research Use Only. Not for use in diagnostic procedures.

Product Information

Contents: Anti-Human CD38 Purified

 **Catalog Number:** 14-0389

Clone: HIT2

Concentration: 0.5 mg/mL

Host/Isotype: Mouse IgG1, kappa

HLDA Workshop: III 155

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C.



Batch Code: Refer to Vial



Use By: Refer to Vial



Contains sodium azide

Description

The HIT2 monoclonal antibody reacts with the human CD38 molecule, an approximately 45 kDa type II transmembrane protein. CD38 is expressed by thymocytes, peripheral B cells including plasma cells, activated T cells, and monocytes. CD38 is a counter-receptor for CD31 and has both cyclase and hydrolase enzymatic activity.

Applications Reported

The HIT2 antibody has been reported for use in flow cytometric analysis, and immunohistochemical staining.

Applications Tested

The HIT2 antibody has been tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at less than or equal to 1 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Horenstein AL, Stockinger H, Imhof BA, Malavasi F. CD38 binding to human myeloid cells is mediated by mouse and human CD31. *Biochem J.* 1998 Mar 15;330 (Pt 3):1129-35.

Deaglio S, Morra M, Mallone R, Ausiello CM, Prager E, Garbarino G, Dianzani U, Stockinger H, Malavasi F. Human CD38 (ADP-ribosyl cyclase) is a counter-receptor of CD31, an Ig superfamily member. *J Immunol.* 1998 Jan 1;160(1):395-402.

Joshua D, Petersen A, Brown R, Pope B, Snowdon L, Gibson J. The labelling index of primitive plasma cells determines the clinical behaviour of patients with myelomatosis. *Br J Haematol.* 1996 Jul;94(1):76-81.

McMichael, AJ, et al. eds. (1987) *Leucocyte Typing III: White Cell Differentiation Antigens.* Oxford University Press. New York.

Related Products

11-4011 Anti-Mouse IgG FITC

14-4714 Mouse IgG1 K Isotype Control Purified (P3.6.2.1)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com