


## Anti-Mouse IL-21 Receptor Purified

Catalog Number: 14-1219

Also Known As: Interleukin-21 Receptor, IL-21R

RUO: For Research Use Only



Staining of 2-day Anti-Mouse CD3e Functional Grade Purified (cat 16-0031) stimulated BALB/c splenocytes with Anti-Mouse CD4 APC (cat. 17-0041) and 0.25 µg of Rat IgG2a κ Isotype Control Purified (cat. 14-4321) (left) or 0.25 µg of Anti-Mouse IL-21 Receptor Purified (right) followed by Anti-Rat IgG Biotin (cat. 13-4813) and then Streptavidin PE (cat. 12-4317). Total viable cells were used for analysis.

### Product Information

Contents: Anti-Mouse IL-21 Receptor Purified


**REF** Catalog Number: 14-1219

Clone: eBio4A9

Concentration: 0.5 mg/ml


Host/Isotype: Rat IgG2a, κ

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C.

**LOT** Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

### Description

The IL-21 receptor (IL-21R) combines with the common γ-chain to form a functional receptor for IL-21. IL-21R is expressed on several lineages including both resting and activated B, T, natural killer and dendritic cells. Binding of its ligand, IL-21, in these cells results in the activation of the Jak/Stat signal transduction pathway. The biological effects of ligand binding are diverse and depend on the cellular context and differentiation state of the cell. In the mouse, IL-21 binding to IL-21R induces the apoptosis of resting and activated B cells, enhances anti-CD3-mediated proliferation of T cells, and inhibits the maturation of dendritic cells and the IL-15-mediated expansion of NK cells.

### Applications Reported

This eBio4A9 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

This eBio4A9 antibody has been tested by flow cytometric analysis of CD3 stimulated mouse splenocytes. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Jin H, Carrio R, Yu A, Malek TR. Distinct activation signals determine whether IL-21 induces B cell costimulation, growth arrest, or Bim-dependent apoptosis. *J Immunol.* 2004 Jul 1;173(1):657-65.

Vosshenrich CA, Ranson T, Samson SI, Corcuff E, Colucci F, Rosmaraki EE, Di Santo JP. Roles for common cytokine receptor gamma-chain-dependent cytokines in the generation, differentiation, and maturation of NK cell precursors and peripheral NK cells in vivo. *J Immunol.* 2005 Feb 1;174(3):1213-21.

### Related Products

11-4317 Streptavidin FITC

11-4811 Anti-Rat IgG FITC

12-4317 Streptavidin PE

13-4813 Anti-Rat IgG Biotin (Polyclonal)

14-4321 Rat IgG2a K Isotype Control Purified

17-4317 Streptavidin APC

---

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.eBioscience.com](http://www.eBioscience.com) • [info@eBioscience.com](mailto:info@eBioscience.com)