

An Affymetrix Company

Anti-Mouse AIRE Purified

Catalog Number: 14-5934

Also known as: Autoimmune Regulator

RUO: For Research Use Only. Not for use in diagnostic procedures.

Product Information

Contents: Anti-Mouse AIRE Purified

REF Catalog Number: 14-5934

Clone: 5H12

Concentration: 0.5 mg/mL Host/Isotype: Rat IgG2c, kappa **Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer **Temperature Limitation:** Store at 2-8°C.

Batch Code: Refer to vial Use By: Refer to vial Contains sodium azide



Description

This 5H12 monoclonal antibody reacts with mouse Autoimmune Regulator (AIRE). This 58-kDa transcription factor is highly expressed within the perinuclear region of mature medullary thymic epithelial cells (mTECs). However, expression of AIRE protein in the periphery remains unclear despite the detection of AIRE mRNA in the lymph node, lymphoid stromal cells, monocytes, and dendritic cells. Studies demonstrate that AIRE is involved in thymocyte development, in particular the regulation of tissue-specific antigen expression during negative selection. Other biological functions for AIRE include chemokine expression and antigen presentation.

The 5H12 antibody recognizes mouse AIRE isoforms 1 and 2.

Applications Reported

This 5H12 antibody has been reported for use in flow cytometric analysis, immunoprecipitation, western blotting, and immunohistochemical staining of frozen tissue sections.

Applications Tested

This 5H12 antibody has been tested by immunohistochemistry on frozen 5-week-old mouse thymus and can be used at less than or equal to 5 ug/mL. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Anderson MS, Su MA. Aire and T cell development. Curr Opin Immunol. 2010; 23:1-9.

Hubert FX, Kinkel SA, Webster KE, Cannon P, Crewther PE, Proeitto AI, Wu L, Heath WR, Scott HS. A specific anti-Aire antibody reveals aire expression is restricted to medullary thymic epithelial cells and not expressed in periphery. J Immunol. 2008 Mar 15;180(6):3824-32. (5H12, FC, IHC-frozen, WB)

Rossi SW, Kim MY, Leibbrandt A, Parnell SM, Jenkinson WE, Glanville SH, McConnell FM, Scott HS, Penninger JM, Jenkinson EJ, Lane PJ, Anderson G. RANK signals from CD4(+)3(-) inducer cells regulate development of Aire-expressing epithelial cells in the thymic medulla. J Exp Med. 2007 Jun 11;204(6):1267-72. (5H12, IHC-frozen)

Related Products

26-4826 Anti-Rat IgG TRITC (polyclonal)