


## Anti-Mouse CD270 (HVEM) Purified

Catalog Number: 14-5962

Also Known As: herpes virus entry mediator, TNFRSF14, LIGHTR

RUO: For Research Use Only



Staining of BALB/c splenocytes with 0.125 µg of Armenian Hamster IgG Isotype Control Purified (cat. 14-4888) (open histogram) or 0.125 µg of Anti-Mouse CD270 (HVEM) Purified (filled histogram) followed by Anti-Armenian Hamster IgG PE (cat. 12-4112). CD3+ cells in the lymphocyte gate were used for analysis.

### Product Information

Contents: Anti-Mouse CD270 (HVEM) Purified

**REF** Catalog Number: 14-5962

Clone: LH1

Concentration: 0.5 mg/ml

Host/Isotype: Armenian Hamster IgG

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer



Temperature Limitation: Store at 2-8°C.



Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

### Description

The LH1 monoclonal antibody reacts with mouse Herpes Virus Entry Mediator (HVEM, TR2), a member of the TNF-receptor superfamily. HVEM is found on most cell types, including T cells, B cells, monocytes, neutrophils and dendritic cells. This receptor was identified as a cellular mediator of herpes simplex virus (HSV) entry. Binding of HSV viral envelope glycoprotein D (gD) to this receptor protein has been shown to be part of the viral entry mechanism. The cytoplasmic region of HVEM was found to bind to several TRAF family members, which may mediate the signal transduction pathways that activate the immune response. HVEM has also been demonstrated to be a unique ligand for BTLA (B and T lymphocyte attenuator). The conservation of the BTLA-HVEM interaction between mouse and human suggests that this system is an important pathway regulating lymphocyte activation and/or homeostasis in the immune response. The LH1 antibody has been reported as a blocking antibody, interfering with the HVEM-LIGHT interaction but not the HVEM-BTLA interaction.

### Applications Reported

This LH1 antibody has been reported for use in flow cytometric analysis and functional assays.

### Applications Tested

This LH1 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Anand S, Wang P, Yoshimura K, Choi IH, Hilliard A, Chen YH, Wang CR, Schulick R, Flies AS, Flies DB, Zhu G, Xu Y, Pardoll DM, Chen L, Tamada K. Essential role of TNF family molecule LIGHT as a cytokine in the pathogenesis of hepatitis. *J Clin Invest.* 2006 Apr;116(4):1045-51. (LH1, FA, PubMed)

Fan Z, Yu P, Wang Y, Wang Y, Fu ML, Liu W, Sun Y, Fu YX. NK-cell activation by LIGHT triggers tumor-specific CD8+ T-cell immunity to reject established tumors. *Blood.* 2006 Feb 15;107(4):1342-51. (PubMed)

Granger SW, Rickert S. LIGHT-HVEM signaling and the regulation of T cell-mediated immunity. *Cytokine Growth Factor Rev.* 2003 Jun-Aug;14(3-4):289-96. (PubMed)

### Related Products

11-4111 Anti-Armenian Hamster IgG FITC

11-4317 Streptavidin FITC

12-4112 Anti-Armenian Hamster IgG PE (Polyclonal)

12-4317 Streptavidin PE  
13-4113 Anti-Armenian Hamster IgG Biotin (Polyclonal)  
14-4888 Armenian Hamster IgG Isotype Control Purified (eBio299Arm)  
14-5969 Anti-Human CD270 (HVEM) Purified (eBioHVEM-122)  
17-4317 Streptavidin APC

---

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.eBioscience.com](http://www.eBioscience.com) • [info@eBioscience.com](mailto:info@eBioscience.com)