


Anti-Mouse CD96 PE

Catalog Number: 12-0960

Also known as: TACTILE

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57Bl/6 splenocytes with Anti-Mouse CD96 PE (cat. 12-0960) and 0.25 ug of Rat IgG2a kappa Isotype Control PE (cat. 12-4321) (left) or 0.25 ug of Anti-Mouse CD96 PE (right). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Mouse CD96 PE
Catalog Number: 12-0960
Clone: 6A6
Concentration: 0.2 mg/mL
Host/Isotype: Rat IgG2a, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer
Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material.
Batch Code: Refer to vial
Use By: Refer to vial

Description

This 6A6 monoclonal antibody reacts with mouse CD96, a cell adhesion receptor also known as TACTILE. A member of the Ig superfamily, CD96 is a single pass transmembrane protein consisting of three extracellular Ig-like domains and a proline/serine/threonine rich region. CD96 is expressed on resident monocytes and T cells at all developmental stages (including $\gamma\delta$ T cells, thymocytes, and NK cells). In the mouse, this receptor is not expressed on B cells. Unlike in humans, mouse CD96 does not exist as alternative splice variants. Reports indicate that CD96 binds CD155/poliovirus receptor and Nectin-1 to mediate NK cell adhesion to target cells and cytotoxicity.

This monoclonal antibody has been reported to block CD96 interaction with CD155 and Nectin-1.

Applications Reported

This 6A6 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This 6A6 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.5 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Meyer D, Seth S, Albrecht J, Maier MK, du Pasquier L, Ravens I, Dreyer L, Burger R, Gramatzki M, Schwinzer R, Kremmer E, Foerster R, Bernhardt G. CD96 interaction with CD155 via its first Ig-like domain is modulated by alternative splicing or mutations in distal Ig-like domains. J Biol Chem. 2009 Jan 23;284(4):2235-44.

Seth S, Maier MK, Qiu Q, Ravens I, Kremmer E, Förster R, Bernhardt G. The murine pan T cell marker CD96 is an adhesion receptor for CD155 and nectin-1. Biochem Biophys Res Commun. 2007 Dec 28;364(4):959-65. (6A6, FC, Pubmed)

Fuchs A, Cella M, Giurisato E, Shaw AS, Colonna M. Cutting edge: CD96 (tactile) promotes NK cell-target cell

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com

Anti-Mouse CD96 PE

Catalog Number: 12-0960

Also known as: TACTILE

RUO: For Research Use Only. Not for use in diagnostic procedures.

adhesion by interacting with the poliovirus receptor (CD155). J Immunol. 2004 Apr 1;172(7):3994-8.

Related Products

12-4321 Rat IgG2a K Isotype Control PE (eBR2a)

17-0032 Anti-Mouse CD3 APC (17A2)

17-1551 Anti-Mouse CD155 APC (TX56)