

Product Data Sheet

FITC anti-human/mouse CD49f

Catalog # / Size: 313605 / 25 tests
313606 / 100 tests

Clone: GoH3

Isotype: Rat IgG2a, κ


Immunogen: Mouse mammary tumor cells

Reactivity: Human, Mouse, **Cross-Reactivity:** Baboon, Chimpanzee, Capuchin Monkey, Cynomolgus, Rhesus, Horse (Equine), Cattle (Bovine, Cow), Sheep (Ovine), Swine (Pig, Porcine), Dog (Canine), Cat (Feline), Rabbit (Lapine)

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with GoH3 FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation^{1,5}, *in vitro* and *in vivo* blocking of cell binding to laminin and blocking the function of integrin α_6 ^{1,4}, and immunohistochemistry of acetone-fixed frozen sections^{2,3,5}. The GoH3 antibody has been reported to block laminin binding *in vitro* and to block integrin α_6 function *in vivo*.

Application References:

- Georas SN, *et al.* 1993. *Blood* 82:2872. (IP, Block)
- Honda T, *et al.* 1995. *J. Clin. Endocrinol. Metab.* 80:2899. (IHC)
- Sonnenberg A, *et al.* 1986. *J. Histochem. Cytochem.* 34:1037. (IHC)
- Nakamura K, *et al.* 1997. *Biochem. Biophys. Res. Commun.* 235:524. (Block)
- Sonnenberg A, *et al.* 1987. *J. Biol. Chem.* 262:10376. (IP, IHC)
- Deregibus MC, *et al.* 2007. *Blood* doi:10.1182/blood-2007-03-078709.
- Horwitz KB, *et al.* 2008. *Proc Natl Acad Sci USA.* 105:5774. PubMed
- Nardella C, *et al.* 2009. *Sci Signal.* 2:55. PubMed
- Xu T, *et al.* 2010. *Mol Cancer Ther.* 9:438. PubMed
- Stepp MA, *et al.* 2007. *J Cell Sci.* 120:2851. PubMed
- Jo M, *et al.* 2010. *Cancer Res.* 70:8948. PubMed
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Grange C, *et al.* 2011. *Cancer Res.* 71:5346. PubMed
- Lai KP, *et al.* 2012. *Mol Endocrinol.* 26:52. PubMed
- Oeztuerk-Winder F, *et al.* 2012. *EMBO J.* 31:3431. (FC) PubMed

Description: CD49f is a 120 kD integrin family member also known as VLA-6 α chain and α_6 integrin subunit. CD49f associates with either integrin β_1 (CD29) or integrin β_4 (CD104) to form receptors (VLA-6 or $\alpha_6\beta_4$ complex) for laminin and kalinin. CD49f is expressed on platelets, monocytes, T cells, placental trophoblasts, epithelial and endothelial cells. CD49f is involved in adhesion and can act as a co-stimulatory molecule for T cell activation and proliferation.

Antigen References:

- Sonnenberg A, *et al.* 1990. *J. Cell Biol.* 110:2145.
- Sonnenberg A, *et al.* 1990. *J. Cell. Sci.* 96:207.
- Aumailley M, *et al.* 1990. *Exp. Cell Res.* 188:55.
- Niessen CM, *et al.* 1994. *Exp. Cell Res.* 211:360.

Related Products:

Product
Cell Staining Buffer
RBC Lysis Buffer (10X)
FITC Rat IgG2a, κ Isotype Ctrl
Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone

RTK2758

Application

FC, ICC, ICFC
FC, ICFC
FC, ICFC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.