


## Anti-Human CD243 (ABCB1) PE

**Catalog Number:** 12-2439

**Also known as:** P-glycoprotein-1, MDR1

**RUO: For Research Use Only. Not for use in diagnostic procedures.**



Staining of the 8226/S (left) and 8226 DOX40 (right) cell lines with Mouse IgG2a K Isotype Control PE (cat. 12-4724) (blue histogram) or Anti-Human CD243 (ABCB1) PE (purple histogram). Total viable cells were used for analysis.

### Product Information



**Contents:** Anti-Human CD243 (ABCB1) PE

**Catalog Number:** 12-2439

**Clone:** UIC2

**Concentration:** 5  $\mu$ L (0.5  $\mu$ g)/test

**Host/Isotype:** Mouse IgG2a



**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

**Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

**Batch Code:** Refer to vial



**Use By:** Refer to vial

**Caution, contains Azide**

### Description

This monoclonal antibody reacts with human Multidrug Resistant (MDR)-1, which is also known as P-glycoprotein (Pgp) and CD243. A 170-kDa transmembrane protein, MDR-1 is an ATP-dependent efflux pump for lipophilic compounds, including anti-cancer drugs. Expression of MDR-1 has been shown to correlate with multidrug resistance. In fact, tumor resistance to chemotherapy has been linked to MDR-1 expression in many cancers. MDR-1 is expressed in a variety of tissues, including the brain, kidney, liver, pancreas, and testes. Within the immune system, this molecule can be found on normal T, B, and natural killer cells, but not on monocytes.

This antibody has been reported to inhibit MDR1-mediated efflux.

### Applications Reported

This UIC2 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

This UIC2 antibody has been pre-titrated and tested by flow cytometric analysis on multidrug resistant cell lines. This can be used at 5  $\mu$ L (0.5  $\mu$ g) per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from  $10^5$  to  $10^8$  cells/test.

### References

Chaudhary PM, Mechetner EB, Roninson IB. Expression and activity of the multidrug resistance P-glycoprotein in human peripheral blood lymphocytes. *Blood*. 1992 Dec 1;80(11):2735-9. (UIC2, FC)

Mechetner EB, Roninson IB. Efficient inhibition of P-glycoprotein-mediated multidrug resistance with a monoclonal antibody. *Proc Natl Acad Sci U S A*. 1992 Jul 1;89(13):5824-8. (UIC2, FA)

### Related Products

12-4724 Mouse IgG2a K Isotype Control PE

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.ebioscience.com](http://www.ebioscience.com) • [info@ebioscience.com](mailto:info@ebioscience.com)

---

## Anti-Human CD243 (ABCB1) PE

**Catalog Number:** 12-2439

**Also known as:** P-glycoprotein-1, MDR1

**RUO: For Research Use Only. Not for use in diagnostic procedures.**

---