


Anti-Human Ki-67 PE

Catalog Number: 12-5699

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of unstimulated (left) normal human peripheral blood cells with Anti-Human CD3 PerCP-Cy5.5 (cat. 46-0036) and Anti-Human Ki-67. Normal human peripheral blood cells were also stimulated with PHA then stained with Mouse IgG1 kappa Isotype Control PE (cat. 12-4714) (blue) or Anti-Human Ki-67 PE (purple). All staining was done using the Foxp3 buffers and protocol.

Product Information

Contents: Anti-Human Ki-67 PE
Catalog Number: 12-5699
Clone: 20Raj1
Concentration: 5 μ L (0.06 μ g)/test
Host/Isotype: Mouse IgG1

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer
Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.
Batch Code: Refer to vial
Use By: Refer to vial

Description

The monoclonal antibody 20Raj1 recognizes the human Ki-67 protein. Two isoforms of Ki-67 exist, a 345 and 395 kDa form that are expressed in dividing cells. Ki-67 is expressed in all cell types and is detectable during active phases of the cell cycle (G1, S, G2, and mitosis) but is absent from resting cells (G0). During interphase, Ki-67 expression is localized to the nucleus but redistributes to the chromosomes during mitosis and has specifically been found to associate with heterochromatin-bound proteins such as chromobox protein homolog 3 (CBX3). In studies of tumor cells, Ki-67 expression has been used as a marker for determining the fraction of proliferating cells within a given population of tumor cells.

This monoclonal antibody 20Raj1 recognizes canine Ki-67.

Applications Reported

This 20Raj1 antibody has been reported for use in intracellular staining followed by flow cytometric analysis. Refer to Foxp3 protocol for optimal staining.

Applications Tested

This 20Raj1 antibody has been pre-titrated and tested by intracellular staining using Foxp3 staining buffers (cat. 00-5523) and flow cytometric analysis of normal human peripheral blood cells. This can be used at 5 μ L (0.06 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Jiao Y, Hua W, Zhang T, Zhang Y, Ji Y, Zhang H, Wu H. Characteristics of CD8+ T cell subsets in Chinese patients with chronic HIV infection during initial ART. *AIDS Res Ther.* 2011 Mar 25;8:15. (**20Raj1**, FC, PubMed)

Schlüter C, Duchrow M, Wohlenberg C, Becker MH, Key G, Flad HD, Gerdes J. The cell proliferation-associated antigen of antibody Ki-67: a very large, ubiquitous nuclear protein with numerous repeated elements, representing a new kind of cell cycle-maintaining proteins. *J Cell Biol.* 1993 Nov;123(3):513-22

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com

Anti-Human Ki-67 PE

Catalog Number: 12-5699

RUO: For Research Use Only. Not for use in diagnostic procedures.

Gerdes J, Lemke H, Baisch H, Wacker HH, Schwab U, Stein H. Cell cycle analysis of a cell proliferation-associated human nuclear antigen defined by the monoclonal antibody Ki-67. J Immunol. 1984 Oct;133(4):1710-5.

Related Products

12-4714 Mouse IgG1 K Isotype Control PE (P3.6.2.8.1)

46-0036 Anti-Human CD3 PerCP-eFluor® 710 (SK7)