


Anti-Mouse CD314 (NKG2D) PE

Catalog Number: 12-5882

Also Known As: KLIRK1

RUO: For Research Use Only



Staining of C57BL/6 splenocytes with Anti-Mouse NK1.1 APC (cat. 17-5941) and 0.125 ug of Rat IgG1 K Isotype Control PE (cat. 12-4301) (left) or 0.125ug of Anti-Mouse CD314 (NKG2D) PE (right). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD314 (NKG2D) PE


REF **Catalog Number:** 12-5882

Clone: CX5

Concentration: 0.2 mg/mL


Host/Isotype: Rat IgG1, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 **Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.

LOT **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

 **Caution, contains Azide**

Description

The CX5 monoclonal antibody reacts with the mouse NKG2D, a lectin-like molecule expressed on both human and mouse NK cells. Mouse NKG2D binds to retinoic acid-inducible RAE-1 α , - β , - γ , - δ , - ϵ and the minor histocompatibility molecule H60 and has the ability to costimulate multiple NK activation receptors, through the DAP12/DAP10 adaptor molecules. NKG2D is expressed by all spleen and liver NK cells, NK1.1⁺ thymocytes, *in vitro* activated LAK cells, and a subset of splenic NKT cells.

Applications Reported

The CX5 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The CX5 antibody has been tested by flow cytometric analysis of mouse splenocyte suspensions. This can be used at less than or equal to 0.5 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Lodoen M, Ogasawara K, Hamerman JA, Arase H, Houchins JP, Mocarski ES, Lanier LL. 2003. NKG2D-mediated natural killer cell protection against cytomegalovirus is impaired by viral gp40 modulation of retinoic acid early inducible 1 gene molecules. *J Exp Med.* 197(10):1245-53.

Cerwenka A, Baron JL, Lanier LL. 2001. Ectopic expression of retinoic acid early inducible-1 gene (RAE-1) permits natural killer cell-mediated rejection of a MHC class I-bearing tumor *in vivo*. *Proc Natl Acad Sci U S A.* 98(20):11521-6.

Cerwenka A, Bakker AB, McClanahan T, Wagner J, Wu J, Phillips JH, Lanier LL. 2000. Retinoic acid early inducible genes define a ligand family for the activating NKG2D receptor in mice. *Immunity.* 12(6):721-7.

Related Products

12-4301 Rat IgG1 K Isotype Control PE

16-5872 Anti-Mouse CD314 (NKG2D) Functional Grade Purified (A10)

16-5873 Anti-Mouse CD314 (NKG2D) Functional Grade Purified (C7)

17-5941 Anti-Mouse NK1.1 APC (PK136)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com