


## Anti-Mouse B7-H4 PE

Catalog Number: 12-5972

Also Known As: B7H4, B7S1, B7-S1, B7X

RUO: For Research Use Only



Staining of non-transfected (left) and mouse B7-H4-transfected (right) L5178Y cells with 0.25 µg of Rat IgG2b κ Isotype Control PE (cat. 12-4032) (open histogram) or 0.25 µg of Anti-Mouse B7-H4 PE (filled histogram). Total viable cells were used for analysis.

### Product Information

Contents: Anti-Mouse B7-H4 PE


**REF** Catalog Number: 12-5972

Clone: 188

Concentration: 0.2 mg/ml


Host/Isotype: Rat IgG2b, κ

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

**LOT** Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

### Description

The 188 monoclonal antibody was generated against and reacts with mouse B7-H4 also known as B7S1, B7X. Cross reactivity of this antibody to other proteins has not been determined. B7-H4 is a newly discovered member of the B7 family reported to inhibit T cell activation, cell cycle progression and IL-2 production. The ligand for B7-H4 has not been identified yet. Simultaneous double staining of cells with two anti-mouse B7-H4 antibodies, Clone 9 and 188, suggests that epitopes recognized by these mAbs are different and/or there is no steric hindrance when antibodies are used together. 188 stains mouse B7-H4 transfected cells and not spleen cells. Exact expression pattern of B7-H4 has not been fully characterized.

### Applications Reported

The 188 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

The 188 antibody has been tested by flow cytometric analysis of transfected cells. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Prasad DV, Richards S, Mai XM, Dong C. 2003. B7S1, a novel B7 family member that negatively regulates T cell activation. *Immunity*. 18(6):863-73.

Choi IH, Zhu G, Sica GL, Strome SE, Cheville JC, Lau JS, Zhu Y, Flies DB, Tamada K, Chen L. 2003. Genomic organization and expression analysis of B7-H4, an immune inhibitory molecule of the B7 family. *J Immunol*. 171(9):4650-4.

Sica GL, Choi IH, Zhu G, Tamada K, Wang SD, Tamura H, Chapoval AI, Flies DB, Bajorath J, Chen L. 2003. B7-H4, a molecule of the B7 family, negatively regulates T cell immunity. *Immunity*. 18(6):849-61.

Chen Y, Yang C, et al. 2006. Expression of the novel co-stimulatory molecule B7-H4 by renal tubular epithelial cells. *Kidney Int*. 2006 Dec;70(12):2092-9. (FA, PubMed)

### Related Products

12-5970 Anti-Mouse B7-H4 PE (Clone 9)

13-5970 Anti-Mouse B7-H4 Biotin (Clone 9)

---

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.eBioscience.com](http://www.eBioscience.com) • [info@eBioscience.com](mailto:info@eBioscience.com)