

Product Data Sheet

Alexa Fluor® 488 anti-mouse CD127 (IL-7Rα)

Catalog # / Size: 135017 / 25 µg

135018 / 100 µg

Clone: A7R34 **Isotype:** Rat IgG2a, κ

Immunogen: IL-7Ra-IgG1 fusion protein

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with

Alexa Fluor® 488 under optimal conditions. The solution is free of

unconjugated Alexa Fluor® 488.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each

* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at

** Alexa Fluor® 488 is a registered trademark of Molecular Probes, Inc. Alexa Fluor® 488 dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending

and issued patents.

Application Notes: In process development has shown A7R34 is able to block clone SB/199

binding to IL-7R.

Application References: 1. Sudo T, et al. 1993. P. Natl. Acad. Sci. USA 90:9125.

2. Hashi H, et al. 2001. J. Immunol. 166:3702. 3. Taylor R, *et al.* 2007. *J. Immunol.* 178:5659. 4. Mazzon C, *et al.* 2011. *Blood.* 118:2733. PubMed

5. Jin J, et al. 2011. J. Immunol. doi:10.4049/jimmunol.1001238. PubMed

A7R34 Alexa Fluoris488 C57BL/6 mouse splenocytes stained with CD3 (145-2C11) APC and A7R34 Alexa Fluor® 488 (top) or rat IgG2a, κ Alexa Fluor® 488 isotype control (bottom). (145-2C11) APC

rat IgG2a.k Alexa Fluor@488

Description: CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7R α . It forms a heterodimer with the common γ chain (γ c or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.

Antigen References:

Sudo T, et al. 1993. P. Natl. Acad. Sci. USA 90:9125.
Okuno Y, et al. 2001. P. Natl. Acad. Sci. USA 99:6246.

3. Pillai M, et al. 2004. Leukemia Lymphoma 45:2403.

Application Related Products: Product Clone Alexa Fluor® 488 Rat IgG2a, κ Isotype Ctrl RTK2758 FC, ICFC

Cell Staining Buffer FC, ICC, ICFC FC, ICFC RBC Lysis Buffer (10X) TruStain fcX™ (anti-mouse CD16/32) 93



