


An Affymetrix Company

## **Anti-Mouse CD11a APC**

Catalog Number: 17-0111

Also known as: Integrin alpha L, Lymphocyte Function-Associated Antigen RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57Bl/6 splenocytes with 0.25 ug of Rat IgG2a K Isotype Control APC (cat. 17-4321) (blue histogram) or 0.25 ug of Anti-Mouse CD11a APC (purple histogram). Total viable cells were used for analysis.

#### **Product Information**

Contents: Anti-Mouse CD11a APC

REF Catalog Number: 17-0111

**Clone:** M17/4

Concentration: 0.2 mg/mL Host/Isotype: Rat IgG2a, kappa



**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer **Temperature Limitation:** Store at 2-8°C. Do not

freeze. Light-sensitive material. **Batch Code:** Refer to vial



Use By: Refer to vial
Contains sodium azide



### Description

The M17/4 monoclonal antibody reacts with mouse CD11a, the 180 kDa integrin alpha L, also known as the lymphocyte function associated antigen-1 (LFA-1) alpha chain. LFA-1, formed by non-covalent association of CD11a with CD18 (integrin beta 2), serves as an important adhesion molecule involved in lymphocyte and granulocyte function. CD54 (ICAM-1), CD102 (ICAM-2), and CD50 (ICAM-3) are ligands for LFA-1. CD11a is expressed by all leukocytes.

#### **Applications Reported**

This M17/4 antibody has been reported for use in flow cytometric analysis.

#### **Applications Tested**

This M17/4 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to  $0.5~\mu g$  per test. A test is defined as the amount ( $\mu g$ ) of antibody that will stain a cell sample in a final volume of 100  $\mu L$ . Cell number should be determined empirically but can range from  $10^5$  to  $10^8$  cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

#### References

Sanchez-Madrid F, Simon P, Thompson S, Springer TA. Mapping of antigenic and functional epitopes on the alphaand beta-subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. J Exp Med. 1983 Aug 1;158(2):586-602.

Sanchez-Madrid F, Davignon D, Martz E, Springer TA. Sanchez-Madrid F, Davignon D, Martz E, Springer TA. Cell Immunol. 1982 Oct;73(1):1-11.

#### **Related Products**

00-4222 Flow Cytometry Staining Buffer 00-4300 10X RBC Lysis Buffer (Multi-species) 17-4321 Rat IgG2a K Isotype Control APC (eBR2a)



# **Anti-Mouse CD11a APC**

Catalog Number: 17-0111

Also known as: Integrin alpha L, Lymphocyte Function-Associated Antigen RUO: For Research Use Only. Not for use in diagnostic procedures.