


Anti-Human CD14 APC

Catalog Number: 17-0149 RUO: For Research Use Only. Not for use in diagnostic procedures.



Description

The 61D3 monoclonal antibody reacts with human CD14, a 53-55 kDa GPI-linked glycoprotein. CD14 is expressed on monocytes, interfollicular macrophages and some dendritic cells. Complexes of LPS and LBP (LPS-Binding Protein) bind with high affinity to monocytes through the surface CD14.

Applications Reported

The 61D3 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This 61D3 antibody has been pre-titrated and tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at 5 μ L (0.25 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

References

Schlossman, S., L. Bloumsell, et al. eds (1995). Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press. New York.

Kishimoto, T., A.E.G., von dem Borne, et al. eds. (1998). Leucocyte Typing VI: White Cell Differentiation Antigens. Garland Publishing Inc. London.

Fadok VA, Warner ML, Bratton DL, Henson PM. CD36 is required for phagocytosis of apoptotic cells by human macrophages that use either a phosphatidylserine receptor or the vitronectin receptor (alpha v beta 3). J Immunol 1998 Dec 1;161(11):6250-7.

Related Products

17-4714 Mouse IgG1 K Isotype Control APC (P3.6.2.8.1)