

FITC anti-mouse CD200R (OX2R)

Catalog # / Size: 123909 / 50 µg
123910 / 500 µg

Clone: OX-110

Isotype: Rat IgG2a, κ

Immunogen: Fusion protein mouse CD200RCD4d3


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.



C57BL/6 bone marrow cells stained with OX-110 FITC (gated on myeloid cell populated)

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional applications (for the relevant formats) include: *in vivo* studies³. The LEAF™ purified antibody is recommended for functional assays.

Application References: 1. Voulgaraki D, *et al.* 2005. *Immunol.* 115:337
2. Stumper TL, *et al.* 2011. *J. Immunol.* 186:1970. PubMed
3. Talebian F, *et al.* 2012. *PLoS One* 7:e31442. (FA)

Description: CD200R, also known as OX2R, is a membrane glycoprotein with up to 70% of its weight derived from N-linked glycosylation. CD200R is expressed primarily by monocytes and neutrophils but also by other leukocytes including T lymphocytes and mast cells. The interaction between CD200 and CD200R may contribute to pathways that suppress and limit macrophage induced inflammatory damage in tissue. Studies suggest that CD200-CD200R interaction may be involved in the control of myeloid cellular function. Levels of expression on resting peripheral blood cells are relatively low.

Antigen References: 1. Voulgaraki D, *et al.* 2005. *Immunol.* 115:337

Related Products:	Product	Clone	Application
	FITC Rat IgG2a, κ Isotype Ctrl	RTK2758	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.