

Product Data Sheet

PE/Cy7 anti-mouse CD183 (CXCR3)

Catalog # / Size: 126515 / 25 µg
126516 / 100 µg

Clone: CXCR3-173

Isotype: Armenian Hamster IgG

Immunogen: Mouse N-terminus of CXCR3


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 splenocytes stained with CXCR3-173 PE/Cy7 and CD3e (145-2C11) APC


Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone CXCR3-173 does not work for western blot.⁷ The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for blocking (Cat. No. 126517). For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 126526) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed for research use only.



C57BL/6 splenocytes stained with Armenian Hamster IgG isotype PE/Cy7 and CD3e (145-2C11) APC

- Application References:**
1. Krug A, *et al.* 2002. *J. Immunol.* 169:6079.
 2. McAleer JP, *et al.* 2009. *J. Immunol.* 182:5322. PubMed
 3. Sharma R. 2009. *J. Immunol.* 183:3212. (FC) PubMed
 4. Barbi J, *et al.* 2009. *FASEB J.* 23:3990. PubMed
 5. Kimpfler S, *et al.* 2009. *J. Immunol.* 183:6330. PubMed
 6. Dürr C, *et al.* 2010. *Cancer Res.* 70:10170. (Block) PubMed
 7. Uppaluri R, *et al.* 2008. *Transplantation* 86:137.

Description: CD183, also known as CXCR3, is a member of the C-X-C chemokine family, characterized by a pair of cysteine residues separated by a single amino acid. CXCR3 is a 38 kD seven pass transmembrane receptor coupled to G-protein. It mediates Ca²⁺ mobilization and chemotaxis in response to C-X-C chemokines, such as IP10 (CXCL10), MIG (CXCL9), I-TAC (CXCL11) and PF4 (CXCL4). CXCR3 is expressed primarily on activated T lymphocytes, NK cells, and some epithelial cells and endothelial cells. It is not expressed on B cells, monocytes or granulocytes.

Related Products:	Product	Clone	Application
	PE/Cy7 Armenian Hamster IgG Isotype Ctrl	HTK888	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.