

Product Data Sheet

PE anti-mouse CD226 (DNAM-1)

Catalog # / Size: 128805 / 25 µg

128806 / 100 µg

Clone: 10E5

Isotype: Rat IgG2b, κ

Immunogen: Th1 polarized T cell clones

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE under optimal conditions. The solution is free of unconjugated PE and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



C57BL/6 splenocytes stained with CD8 (53-6.7) FITC and 10E5 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For

immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is

recommended that the reagent be titrated for optimal performance for each application.

Application References: 1. Dardalhon V, et al. 2005. J. Immunol. 175:1558

Description: CD226 (DNAM-1) is constitutively expressed on native CD8+ cells and on CD4+ T cells, macrophages and NK

cells. This antibody (10E5) was reported to bind about 40% of inactivated CD4+ cells and binds only to differentiated

Th1 cells, but not to Th2 or Th0 cells. It is also reported to suppress antigen-specific T cell expansion and

EAE(experimental allergic encephalitis) mediated by Th1 cells.

Antigen References: 1.Tahara Hanaoka S, et al. 2006. Blood 107:1491

2. Shibuya K, et al. 2003. J. Exp. Med. 198:1829

Related Products: Product

PE Rat IgG2b, κ Isotype Ctrl

Cell Staining Buffer RBC Lysis Buffer (10X)

TruStain fcX™ (anti-mouse CD16/32)

Clone RTK4530

93

Application FC, ICFC FC, ICC, ICFC

FC, ICFC



