

## **Product Data Sheet**

## Purified anti-mouse CD196 (CCR6)

Catalog # / Size: 129801 / 25 µg

 $129802 / 100 \, \bar{\mu}g$ 

Clone: 29-2L17

Isotype: Armenian Hamster IgG

Immunogen: N-terminal peptide (aa. 2-38) of mouse CCR6 fused with GST

Reactivity: Mouse

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Related Products: Product

**Storage:** The antibody solution should be stored undiluted at 4°C.

## **Applications:**

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Description: CCR6 is a G-protein linked chemokine receptor which binds the chemokine CCL20/MIP-3a. It is expressed on B

lymphocytes and certain subsets of dendritic cells and T cells. CCR6 in reported to be involved in mucosal immune

Clone

**HTK888** 

response and lymphocyte migration and homeostasis.

Antigen References: 1. Rossi D, et al. 1997. J. Immunol. 158:1033.

2. Lukacs N, et al. 2001. J. Exp. Med. 194:551.


3. Cook DN, et al. 2000. Immunity 12:495

Purified Armenian Hamster IgG Isotype Ctrl

Cell Staining Buffer

RBC Lysis Buffer (10X)

C57BL/6 splenocytes stained with purified 29-2L17, followed by biotinylated anti-Armenian hamster IgG and Sav-PE



104



FC, ICC, ICFC, IF, IP, WB FC, ICC, ICFC FC, ICFC



