

Product Data Sheet

PE anti-mouse CD226 (DNAM-1)

Catalog # / Size: 132005 / 25 μg

132006 / 100 µg

Clone: 480.1

Isotype: Rat IgG2a, κ

Immunogen: Jurkat cells transfected with mouse CD226

Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE under optimal conditions. The solution is free of unconjugated PE and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



C57BL/6 splenocytes stained with 53-6.7 (CD8) FITC and 480.1 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For

immunofluorescent staining, the suggested use of this reagent is ≤0.25 μg per million cells in 100 μl volume. It is

recommended that the reagent be titrated for optimal performance for each application.

Application References: Dardalhon V, et al. 2005. J. Immunol. 175:1558

Description: CD226 (DNAM-1) is constitutively expressed on native CD8+ cells and on some CD4+ T cells, macrophages and NK

cells. CD226 is involved in NK and T cell mediated cytotoxicity against certain tumors. CD155 and CD112 are the ligands for CD226. This antibody (480.1) is reported by the developer to partially block the binding of mouse CD155.

93

Antigen References: 1. Tahara Hanaoka S et al. 2006 Blood 107:1491

2. Shibuya K et al. 2003. J. Exp. Med. 198:1829

Related Products: Product Clone Application

PE Rat IgG2a, κ Isotype Ctrl Cell Staining Buffer RBC Lysis Buffer (10X)

TruStain fcX™ (anti-mouse CD16/32)

RTK2758 FC, ICFC

FC, ICFC FC, ICC, ICFC

FC ICFC

FC, IC



