

# Product Data Sheet

## FITC anti-mouse CD244.2 (2B4 B6 Alloantigen)

**Catalog # / Size:** 133503 / 50 µg  
133504 / 500 µg

**Clone:** m2B4 (B6)458.1

**Isotype:** Mouse IgG1, κ

**Immunogen:** m2B4-Fc


**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 splenocytes stained with m2B4(B6)458.1 FITC and NK-1.1 APC

## Applications:

**Applications:** FC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is  $\leq 0.25 \mu\text{g}$  per  $10^6$  cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application References:**

1. Lee KM, et al. 2006. *Blood* Apr 15:107:3181.
2. Schatzle JD, et al. 1999. *P. Natl Acad Sci USA* 96:3870.
3. Stepp SE, et al. 1999. *Eur. J. Immunol.* 29:2392.
4. de la Luz Sierra M, et al. 2010. *Blood* 115:3970. PubMed

**Description:** Mouse CD244, also known as 2B4, is a receptor belonging to the CD2 family of proteins involved in non-MHC-restricted cytotoxicity. It is expressed on all natural killer (NK) cells, IL-2 activated NK (LAK) cells, NKT cells, and a subset of T lymphocytes, including dendritic epidermal T cells. At least there are two isoforms of CD244.2 proteins that differ in the length of the cytoplasmic domain. The long form functions in an inhibitory manner, while the short form functions in an activating manner. The ligand of CD244 is CD48, which expressed on all hematopoietic cells. It was reported that CD244 interacts with CD48 is essential for IL-2-driven expansion and activation of murine NK cells. The 2B4 antibody reacts with CD244.2, the 2B4 alloantigen which is expressed on C57BL/6 mice, but not on Balb/c mice.

**Antigen References:**

1. Brown MH, et al. 1998. *J. Exp. Med.* 188:2083.
2. Davis SJ, et al. 1996. *Immunol Today.* 17:177.
3. Kumaresan PR, et al. 2000. *Immunogenetics.* 51:306.
4. Latchman Y, et al. 1998. *J. Immunol.* 161:5809.

### Related Products:

Product  
 FITC Mouse IgG1, κ Isotype Ctrl  
 Cell Staining Buffer  
 RBC Lysis Buffer (10X)  
 TruStain fcX™ (anti-mouse CD16/32)

Clone  
 MOPC-21

93

Application  
 FC, ICFC  
 FC, ICC, ICFC  
 FC, ICFC  
 FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biollegend.com/ordering#license](http://www.biollegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.