

PE anti-mouse CD272 (BTLA)

Catalog # / Size: 134803 / 25 µg
134804 / 100 µg

Clone: 8F4

Isotype: Mouse IgG1, κ

Immunogen: C57BL/6 BTLA Ig domain protein in CFA


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.


Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 splenocytes stained with 8F4 PE and CD45R/B220 APC



C57BL/6 splenocytes stained with mouse IgG1 PE isotype control and CD45R/B220 APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References: 1. Hurchla MA, *et al.* 2005. *J. Immunol.* 174:3377

Description: B and T lymphocyte attenuator (BTLA) is an Ig superfamily coinhibitory receptor with structural similarity to programmed cell death 1 (PD-1) and CTLA-4. BTLA is expressed on B cells, T cells, macrophages, dendritic cells, NKT cells, and NK cells. Engagement of BTLA by its ligand Herpes Virus Entry Mediator (HVEM) is critical for negatively regulating immune response. The absence of BTLA with HVEM inhibitory interactions leads to increased experimental autoimmune encephalomyelitis severity, enhanced rejection of partially mismatched allografts, an increased CD8⁺ memory T cell population, increased severity of colitis, reduced effectiveness of T regulatory cells. BTLA takes an important role in the induction of peripheral tolerance of both CD4⁺ and CD8⁺ T cells in vivo. Tolerant T cells have significant up-regulated expression of BTLA compared with effector and naïve T cells. BTLA may cooperate with CTLA-4 and PD-1 to control T cell tolerance and autoimmunity. It was reported that BTLA may regulate T cell function by binding to B7-H4. But further studies are needed to confirm. The existence of three distinct BTLA alleles was reported. The BTLA antibody reacts with mouse BTLA from both BALB/c and C57BL/6 strains.

Antigen References:

1. Liu X, *et al.* 2009. *J. Immunol.* 182:4516
2. Miller ML, *et al.* 2009. *J. Immunol.* 183:32
3. Sun Y, *et al.* 2009. *J. Immunol.* 183:1946
4. Vendel AC, *et al.* 2009. *J. Immunol.* 182:1509
5. Watanabe N, *et al.* 2003. *Nat. Immunol.* 4(7):670
6. Sedy JR, *et al.* 2005. *Nat Immunol.* 6(1):90

Related Products:

PE Mouse IgG1, κ Isotype Ctrl	Clone	MOPC-21
Cell Staining Buffer		
RBC Lysis Buffer (10X)		
TruStain fcX™ (anti-mouse CD16/32)	Clone	93

Application

FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.