

APC anti-mouse CD270 (HVEM)

Catalog # / Size: 136305 / 25 µg
136306 / 100 µg

Clone: HMHV-1B18

Isotype: Armenian hamster IgG

Immunogen: Mouse HVEM-hIgG1Fc fusion protein


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 splenocytes stained with HMHV-1B18 APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Description: Herpes Virus Entry Mediator (HVEM, TR2) is a type I transmembrane protein of TNF-receptor superfamily. This receptor was identified as a cellular mediator of herpes simplex virus (HSV) entry. Binding of HSV viral envelope glycoprotein D to this receptor has been shown to be part of the viral entry mechanism. It is expressed on most cell types, including T cells, B cells, monocytes, neutrophils, and dendritic cells. It is also found in brain, heart, kidney, liver, and other organs. The ligands of HVEM are LIGHT, BTLA, LT α , and CD160. HVEM activates NF- κ B through the TNF-related cytokine LIGHT to serve as a costimulatory pathway during T cell activation. HVEM also functions as a ligand for the Ig superfamily members B and T lymphocyte attenuator (BTLA) and CD160 to deliver a coinhibitory signal and limit inflammatory responses initiated by T cells. HVEM plays an important role in regulating lymphocyte activation and homeostasis in immune responses.

- Antigen References:**
1. Fan Z, *et al.* 2006. *Blood* 107:1342
 2. Granger SW, *et al.* 2008. *Stem Cells*. 27:653
 3. Kwon BS, *et al.* 1997. *J. Biol. Chem.* 272:13471
 4. Gonzalez LC, *et al.* 2005. *Proc. Nat. Acad. Sci. U. S. A.* 102:1116
 5. Montgomery RI, *et al.* 1996. *Cell*. 87:427
 6. Cheung TC, *et al.* 2009. *Proc. Nat. Acad. Sci. U. S. A.* 106:6244
 7. Cai G, *et al.* 2009. *Immunol. Rev.* 229:244

Related Products:	Product	Clone	Application
	APC Armenian Hamster IgG Isotype Ctrl	HTK888	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.