

APC anti-human/mouse/rat CD278 (ICOS)

Catalog # / Size: 313509 / 25 µg
313510 / 100 µg

Clone: C398.4A

Isotype: Armenian Hamster IgG

Immunogen: Mouse T cell clone D10.G4.1


Reactivity: Human, Mouse, Rat, **Cross-Reactivity:** Rhesus, Swine (Pig, Porcine)

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



PHA-stimulated human peripheral blood lymphocytes (3 days) stained with C398.4A APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10^6 cells in 100 µl volume. It is recommended that reagents be titrated for optimal performance in the particular application.

Application Notes: The C398.4A antibody is useful for flow cytometric analysis and is able to costimulate T cell activation and proliferation. Additional reported applications (for the relevant formats) include: immunoprecipitation¹, immunohistochemical staining of acetone-fixed frozen sections, and *in vitro* costimulation of T cell activation^{1,3,4}. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 313512).

Application References:

1. Redoglia V, *et al.* 1996. *Eur. J. Immunol.* 26:2781. (FC IP Costim)
2. Yagi J, *et al.* 2003. *J. Immunol.* 171:783. (FC)
3. Arimura Y, *et al.* 2002. *Int. Immunol.* 14:555. (Costim)
4. Arimura Y, *et al.* 2004. *J. Biol. Chem.* 279:11408. (Costim)

Description: ICOS, also known as inducible costimulatory molecule and H4, is a 47-57 kD protein. This protein is homologous to the CD28/CTLA-4 proteins. ICOS is expressed on activated T cells and a subset of thymocytes. It is able to costimulate T cells proliferation. In addition, ICOS is involved in humoral immune responses (B cell germinal center formation). The ICOS ligand is B7h/B7RP-1 or B7-H2. ICOS stimulation has been shown to potentiate TCR-mediated IL-4 and IL-10 production and has been proposed to play a role in Th2 cell development.

Antigen References:

1. Redoglia V, *et al.* 1996. *Eur. J. Immunol.* 26:2781.
2. Hutloff A, *et al.* 1999. *Nature* 397:263.
3. Buonfiglio D, *et al.* 2000. *Eur. J. Immunol.* 30:3463.
4. Coyle AJ, *et al.* 2000. *Immunity* 13:95.

Related Products:Product

Cell Staining Buffer
 APC Armenian Hamster IgG Isotype Ctrl
 Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone

HTK888

Application

FC, ICC, ICFC
 FC, ICFC
 FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.