


## Anti-Mouse CD335 (NKp46) PerCP-eFluor® 710

**Catalog Number:** 46-3351

**Also known as:** NCR1, Ly-94

**RUO: For Research Use Only. Not for use in diagnostic procedures.**



Staining of BALB/c splenocytes with Anti-Mouse CD49b (Integrin alpha 2) PE (cat. 12-5971) and 0.125 ug of Rat IgG2a K Isotype Control PerCP-eFluor® 710 (cat. 46-4321) (left) or 0.125 ug of Anti-Mouse CD335 (NKp46) PerCP-eFluor® 710 (right). Cells in the lymphocyte gate were used for analysis.

### Product Information

**Contents:** Anti-Mouse CD335 (NKp46)  
PerCP-eFluor® 710



**Catalog Number:** 46-3351

**Clone:** 29A1.4

**Concentration:** 0.2 mg/mL

**Host/Isotype:** Rat IgG2a, kappa



**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

**Temperature Limitation:** Store at 2-8°C. Do not freeze. Light sensitive material.



**Batch Code:** Refer to vial



**Use By:** Refer to vial



**Caution, contains Azide**

### Description

The monoclonal antibody 29A1.4 recognizes mouse NKp46, also known as CD335. CD335, a member of the natural cytotoxicity receptor (NCR) family, is a glycoprotein with 2 Ig-like domains and a short cytoplasmic tail. Expression of CD335 is uniquely found on NK cells (including immature NK cells, defined as DX5- CD3-, and thereby allowing discrimination between NKT cells and NK cells (NKp46+, CD3-). Furthermore, unlike many of the NK markers which also stain NKT cells, staining with 29A1.4 is not strain specific. Staining has been shown on C57Bl/6, SJL, CBA/CA and BALB/C strains. NKp46 has been shown to play a role in NK cell-mediated lysis of several tumor cells and pathogen-infected cell lines.

The 29A1.4 monoclonal antibody has been shown to activate NK cells *in vitro*. The 29A1.4 monoclonal antibody does not deplete NK cells *in vivo*.

### Applications Reported

This 29A1.4 antibody has been reported for use in flow cytometric analysis.

### Applications Tested

This 29A1.4 antibody has been tested by flow cytometric analysis of mouse spleen cells. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

PerCP-eFluor® 710 can be used in place of PE-Cy5, PE-Cy5.5 or PerCP-Cy5.5. PerCP-eFluor® 710 emits at 710 nm and is excited with the blue laser (488 nm). Please make sure that your instrument is capable of detecting this fluorochrome. For a filter configuration, we recommend using the 685 LP dichroic mirror and 710/40 band pass filter, however the 695/40 band pass filter is an acceptable alternative.

Our testing indicates that PerCP-eFluor® 710 conjugated antibodies are stable when stained samples are exposed to freshly prepared 2% formaldehyde overnight at 4°C, but please evaluate for alternative fixation protocols.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.ebioscience.com](http://www.ebioscience.com) •  
[info@ebioscience.com](mailto:info@ebioscience.com)

---

## **Anti-Mouse CD335 (NKp46) PerCP-eFluor® 710**

**Catalog Number:** 46-3351

**Also known as:** NCR1, Ly-94

**RUO: For Research Use Only. Not for use in diagnostic procedures.**

---

Click here or contact eBioscience Technical Support for more information on eFluor® Organic Dyes including PerCP-eFluor® 710.

### **References**

Ota N, Wong K, Valdez PA, Zheng Y, Crellin NK, Diehl L, Ouyang W. IL-22 bridges the lymphotoxin pathway with the maintenance of colonic lymphoid structures during infection with *Citrobacter rodentium*. *Nat Immunol*. 2011 Aug 28 doi: 10.1038/ni.2089. (29A1.4, FC, PubMed)

Walzer T, Chiossone L, Chaix J, Calver A, Carozzo C, Garrigue-Antar L, Jacques Y, Baratin M, Tomasello E, Vivier E. Natural killer cell trafficking in vivo requires a dedicated sphingosine 1-phosphate receptor. *Nat Immunol*. 2007 Dec;8(12):1337-44. (29A1.4, FC PubMed)

Walzer T, Bléry M, Chaix J, Fuseri N, Chasson L, Robbins SH, Jaeger S, André P, Gauthier L, Daniel L, Chemin K, Morel Y, Dalod M, Imbert J, Pierres M, Moretta A, Romagné F, Vivier E. Identification, activation, and selective in vivo ablation of mouse NK cells via NKp46. *Proc Natl Acad Sci U S A*. 2007 Feb 27;104(9):3384-9. (29A1.4, FC, FA PubMed)

### **Related Products**

12-5971 Anti-Mouse CD49b (Integrin alpha 2) PE (DX5)

46-4321 Rat IgG2a K Isotype Control PerCP-eFluor® 710 (eBR2a)

48-0032 Anti-Mouse CD3 eFluor® 450 (17A2)