


Anti-Human IL-8 PerCP-eFluor® 710

Catalog Number: 46-8088

Also known as: CXCL8, Interleukin-8, NAP-1

RUO: For Research Use Only. Not for use in diagnostic procedures.



Normal human peripheral blood cells cultured 18 hours in the presence of Protein Transport Inhibitor Cocktail (cat. 00-4980) and stained intracellularly with Anti-Human CD14 FITC (cat. 11-0149) and Anti-Human IL-8 PerCP-eFluor® 710. Cells in the monocyte gate were used for analysis.

Product Information

Contents: Anti-Human IL-8 PerCP-eFluor® 710



Catalog Number: 46-8088

Clone: 8CH

Concentration: 5 µL (0.03 µg)/test

Host/Isotype: Mouse IgG1, kappa



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material.



Batch Code: Refer to vial



Use By: Refer to vial

Description

This 8CH monoclonal antibody reacts with human IL-8 (CXCL8), a pro-inflammatory CXC chemokine. It is synthesized as a 99 amino acid precursor protein that is further processed into one of four isoforms, with the most common being 72 or 77 amino acids in length. IL-8(77) is secreted primarily by endothelial cells and is thought to be a less potent neutrophil activator than the other forms. It is present at high levels during fetal development, where it mediates angiogenesis rather than inflammation. The predominant form present in adults is IL-8(72), which is expressed by monocytes, macrophages, epithelial cells, and fibroblasts in response to inflammatory stimuli, environmental stress, and steroid hormones. IL-8(72) is essential for the activation and recruitment of neutrophils to sites of inflammation, and has also been found to influence T cell migration. Signaling occurs through the G-protein coupled receptors CXCR1 or CXCR2. IL-8 transcripts are often upregulated in tumors, and it is associated with tumor angiogenesis and metastasis.

Applications Reported

This 8CH antibody has been reported for use in intracellular staining followed by flow cytometric analysis.

Applications Tested

This 8CH antibody has been pre-titrated and tested by intracellular staining followed by flow cytometric analysis. This can be used at 5 µL (0.03 µg) per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.

PerCP-eFluor® 710 can be used in place of PE-Cy5, PE-Cy5.5 or PerCP-Cy5.5. PerCP-eFluor® 710 emits at 710 nm and is excited with the blue laser (488 nm). Please make sure that your instrument is capable of detecting this fluorochrome. For a filter configuration, we recommend using the 685 LP dichroic mirror and 710/40 band pass filter, however the 695/40 band pass filter is an acceptable alternative.

Our testing indicates that PerCP-eFluor® 710 conjugated antibodies are stable when stained samples are exposed to freshly prepared 2% formaldehyde overnight at 4°C, but please evaluate for alternative fixation protocols.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Human IL-8 PerCP-eFluor® 710

Catalog Number: 46-8088

Also known as: CXCL8, Interleukin-8, NAP-1

RUO: For Research Use Only. Not for use in diagnostic procedures.

Click [here](#) or contact eBioscience Technical Support for more information on eFluor™ Organic Dyes including PerCP-eFluor® 710.

References

Maheshwari A, Voitenok NN, Akalovich S, Shaik SS, Randolph DA, Sims B, Patel RP, Killingsworth CR, Fallon MB, Ohls RK. Developmental changes in circulating IL-8/CXCL8 isoforms in neonates. *Cytokine*. 2009 Apr;46(1):12-6.

Tajima A, Iwase T, Shinji H, Seki K, Mizunoe Y. Inhibition of endothelial interleukin-8 production and neutrophil transmigration by Staphylococcus aureus beta-hemolysin. *Infect Immun*. 2009 Jan;77(1):327-34.

Waugh DJ, Wilson C. The interleukin-8 pathway in cancer. *Clin Cancer Res*. 2008 Nov 1;14(21):6735-41.

Baggiolini M, Walz A, and Kunkel SL. Neutrophil-activating peptide-1/interleukin 8, a novel cytokine that activates neutrophils. *J Clin Invest*. 1989 Oct;84(4):1045-9.

Related Products

00-4980 Protein Transport Inhibitor Cocktail (500X)

11-0149 Anti-Human CD14 FITC (61D3)

46-4714 Mouse IgG1 K Isotype Control PerCP-eFluor® 710 (P3.6.2.8.1)

88-7087 Human IL-8 ELISA Ready-SET-Go!® (To Be Discontinued. Refer to 2nd Generation RSG Version: cat. 88-8086)