

Product Data Sheet

PE anti-mouse Podoplanin

Catalog # / Size: 127407 / 50 µg
127408 / 200 µg

Clone: 8.1.1

Isotype: Syrian Hamster IgG


Reactivity: Mouse Podoplanin

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



TE-71, mouse thymic epithelial stromal cell line, stained with 8.1.1 PE

Applications:

Applications: FC -Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 µg per 10^6 cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemistry⁶.

Application References:

1. Farr A, *et al.* 1992. *J. Histochem. Cytochem.* 40:651.
2. Farr AG, *et al.* 1992. *J. Exp. Med.* 176:1477.
3. Bekiaris V, *et al.* 2008. *J. Immunol.* 180:6768.
4. Algars A, *et al.* 2011. *Blood* 117:4387. PubMed
5. Reis VO, *et al.* 2012. *Immunobiology.* 217:831. PubMed
6. Kaji C, *et al.* 2012. *Acta. Histochem. Cytochem.* 45:227. (IHC)

Description: The mucin-type glycoprotein podoplanin is thought to be involved in the development of the lymphatic vascular system. Podoplanin is named after its expression in the kidney glomerular epithelial cells (podocytes). It has a potential role in tumor progression.

Antigen References:

1. Farr A, *et al.* 1992. *J. Histochem. Cytochem.* 40:651.
2. Schacht V, *et al.* 2005. *Am. J. Pathol.* 166:913.

Related Products:

Product
Cell Staining Buffer
RBC Lysis Buffer (10X)
PE Syrian Hamster IgG Isotype Ctrl
TruStain fcX™ (anti-mouse CD16/32)

Clone

SHG-1
93

Application

FC, ICC, ICFC
FC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.