

Product Data Sheet

FITC anti-rat CD45

Catalog # / Size: 202205 / 100 µg

Clone: OX-1

Isotype: Mouse IgG1, κ

Immunogen: Enriched glycoprotein fraction from Wistar rat thymocytes


Reactivity: Rat

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.5% BSA (origin U.S.)

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



LOU rat splenocytes stained with OX-1 FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation¹, immunofluorescence microscopy (acetone fixed, ammonium-thiocyanate separated epidermal sheets)², immunohistochemistry of acetone³-, isopentane⁴-, PLP⁵-, and n-hexane⁶ fixed frozen sections and zinc-fixed paraffin-embedded sections, and partial inhibition of NK cell lysis of syngeneic tumor cell lines¹.

Application References:

1. Giezeman-Smits KM, *et al.* 1999. *J. Immunol.* 163:71. (IP)
2. Elbe A, *et al.* 1994. *J. Invest. Dermatol.* 102:74. (IF)
3. Kouwenhoven E, *et al.* 2001. *Kidney Int.* 59:1142. (IHC)
4. Martin A, *et al.* 1995 *Clin. Exp. Immunol.* 22:283. (IHC)
5. Sayegh MH, *et al.* 1995 *J. Exp. Med.* 181:186. (IHC)
6. Morioka Y, *et al.* 2000. *Kidney Int.* 60:2192. (IHC)

Description: CD45 is a 180-220 kD protein also known as leukocyte common antigen (LCA). It is a protein tyrosine phosphatase with multiple isoforms differing as a result of alternative splicing of the extracellular domain and glycosylation. CD45 is expressed on all hematopoietic cells except erythrocytes and platelets; isoform expression depends on cell type, activation state, and cell maturation. CD45 functions in signal transduction through T and B cell antigen receptors. CD45 has been shown to interact with various proteins including galectin-1, CD2, CD3, and CD4. The OX-1 antibody has been shown to partially inhibit NK cell-mediated lysis of syngeneic tumor cells *in vitro*.

Antigen References:

1. Sunderland CA, *et al.* 1979. *Eur. J. Immunol.* 9:155.
2. Woollett GR, *et al.* 1985. *Eur. J. Immunol.* 15:168.

Related Products:

Product
 FITC Mouse IgG1, κ Isotype Ctrl
 Cell Staining Buffer

Clone
 MOPC-21

Application
 FC, ICFC
 FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

