

Product Data Sheet

PerCP anti-human CD1c

Catalog # / Size: 331511 / 25 tests
331512 / 100 tests

Clone: L161

Isotype: Mouse IgG1, κ


Workshop Number: V T-CD01.18

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PerCP under optimal conditions. The solution is free of unconjugated PerCP and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with L161 PerCP

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l volume or 5 μ l per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP has a maximum absorption of 482 nm and 564 nm and a maximum emission of 675 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunocytochemical staining¹.

Application References: 1. M. del Salamone C, *et al.* 2001. *J. Leukoc. Biol.* 70:567.
2. de Fraissinette A, *et al.* 1988. *Exp. Hematol.* 16:764.

Description: CD1c, also known as R7 or M241, is a 43 kD member of the five CD1 antigens (CD1a-e) in human. The CD1 molecules are type I glycoprotein with structural similarities to MHC class I and a non-covalent associated with β_2 -microglobulin, belonging to Ig superfamily. CD1c is expressed on cortical thymocytes, Langerhans cells, dendritic cells, and a subset of B cells. It has been reported that CD1c is also expressed on mature T cells with tightly regulated manner. The function of CD1c is involved in antigen-presentation of glycolipids. It may also act in T cells as an immune regulatory molecule.

Antigen References: 1. Fainboim LM and del C. Salamone. 2002. *J. Biol. Reg. Homeos. Ag.* 16:125.
2. M. del Salamone C, *et al.* 2001. *J. Leukocyte Biol.* 70:567.
3. Zola H, *et al.* Eds. 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers.* P42.

Related Products:	Product	Clone	Application
	PerCP Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.