

Product Data Sheet

PE anti-human CD4

Catalog # / Size: 344605 / 25 tests

344606 / 100 tests

Clone: SK3

Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE under optimal conditions. The solution is free of unconjugated PE and

unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. **Test size products are transitioning from 20 \muI to 5 \muI per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 \muI staining volume or** per 100 µl of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

1. Evans RL, et al. 1981. Immunol. 78:544 Application References:

Arno A et al. 1999. J. Infect. Dis. 180:56
Muech M, et al. 1997. Blood 89:1364

Description: CD4, also known as T4, is a 55 kD single-chain type I transmembrane glycoprotein expressed on most thymocytes, a

subset of T cells, and monocytes/macrophages. CD4, a member of the Ig superfamily, recognizes antigens

associated with MHC class II molecules and participates in cell-cell interactions, thymic differentiation, and signal transduction. CD4 acts as a primary receptor for HIV, binding to HIV gp120. CD4 has also been shown to interact with

Antigen References: 1. Center D et al. 1996. Immunol. Today 17:476.

2. Gaubin M et al. 1996. Eur. J. Clin. Chem. Clin. Biochem. 34:723.

Related Products: Product Clone MOPC-21 PE Mouse IgG1, κ Isotype Ctrl

Cell Staining Buffer RBC Lysis Buffer (10X)

Human TruStain FcX™ (Fc Receptor Blocking Solution)

Relative Cell Number 10⁰ 102 104 103 Log Fluorescence Intensity

Human peripheral blood lymphocytes stained with SK3 PE

Application FC, ICFC FC, ICC, ICFC





