

## PE anti-human CD5

**Catalog # / Size:** 300607 / 25 tests  
300608 / 100 tests

**Clone:** UCHT2

**Isotype:** Mouse IgG1,  $\kappa$


**Workshop Number:** III 518

**Reactivity:** Human, **Cross-Reactivity:** Chimpanzee, Capuchin Monkey, Common Marmoset

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes were stained with CD5 (UCHT2) PE (solid line) or mouse IgG1,  $\kappa$  PE (dashed line).

## Applications:

**Applications:** FC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20  $\mu$ l to 5  $\mu$ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100  $\mu$ l staining volume or per 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at [www.biolegend.com/testsize](http://www.biolegend.com/testsize) regarding the test size change.

**Application Notes:** Additional reported applications (for the relevant formats) include: Western blotting<sup>2</sup> and immunohistochemical staining of acetone-fixed frozen sections<sup>2,5</sup>.

**Application References:**

1. Knapp W, *et al.* 1989. Leucocyte Typing IV Oxford University Press. New York.
2. Renaudineau Y, *et al.* 2005. *Blood* 106:2781. (WB IHC)
3. Porter JC and Hogg N. 1997. *J. Cell Biol.* 138:1437.
4. Saliba AE, *et al.* 2010. *P. Natl. Acad. Sci. USA* 107:14524. PubMed
5. Kap Y, *et al.* 2009. *J. Histochem. Cytochem.* 57:1159. (IHC)

**Description:** CD5 is a 67 kD single chain type I glycoprotein also known as Leu-1, Ly-1 and T1. It is a member of the scavenger receptor superfamily found on T cells, thymocytes, B cell subsets, chronic B lymphocytic leukemia (B-Cells), and peripheral blood dendritic cells. CD5 modulates T and B cell receptor signaling, thymocyte maturation, and T-B cell interactions upon binding to ligands such as CD72.

**Antigen References:**

1. Kipps T. 1988. *Adv. Immunol.* 47:117.
2. Resnick D, *et al.* 1993. *Trends Biochem. Sci.* 19:5.
3. Wood GS, *et al.* 1992. *Am. J. Pathol.* 14:789.

### Related Products: Product

PE anti-human CD19	Clone	Application
PE anti-human CD3	HIB19	FC
PE anti-human CD3	HIT3a	FC
PE Mouse IgG1, $\kappa$ Isotype Ctrl	UCHT1	FC
Cell Staining Buffer	MOPC-21	FC, ICFC
RBC Lysis Buffer (10X)		FC, ICC, ICFC
Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICFC
		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.