

Product Data Sheet

PE anti-human CD43

Catalog # / Size: 343203 / 25 tests
343204 / 100 tests

Clone: CD43-10G7

Isotype: Mouse IgG1, κ


Immunogen: MV4-11 cell line

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**




Human peripheral blood lymphocytes stained with CD43-10G7 PE and CD3 APC (top) or mouse IgG1, κ PE isotype control and CD3 APC (bottom).

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application References: 1. Rosenkranz AR, *et al.* 1993. *Immunology*. 80:431
2. Skubitz KM, *et al.* 1998. *J. Leukoc. Biol.* 64:800



Description: CD43, also known as Ly-48, Leukosialin, Sialophorin, Leukocyte Sialoglycoprotein, and W3/13, is a large single chain of type I transmembrane glycoprotein with abundant O-glycosylation and sialylation sites. It has been reported that CD43 binds to CD54 and Siglec-1. CD43 plays dual roles in cell adhesion and anti-adhesion, as well as costimulation of T cell activation and survival, and induction of apoptosis of T cells and hematopoietic progenitors.

Antigen References: 1. Neil Barclay A, *et al.* 1997. *The Leucocyte Antigen Facts Book: Second Edition*. Page 238
2. Killeen A, *et al.* 1987. *EMBO J.* 6(13):4029.
3. Thurman EC, *et al.* 1998. *Intl. Immunol.* 10:691.
4. Onami TM, *et al.* 2002. *J. Immunol.* 168:6022.
5. Tong J, *et al.* 2004. *J. Exp. Med.* 199:1277.
6. Jones AT, *et al.* 1994. *J. Immunol.* 153:3426.

Related Products:

Product
PE Mouse IgG1, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone
MOPC-21

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.