

Product Data Sheet

PE/Cy5 anti-human CD45RO

Catalog # / Size: 304208 / 100 tests

Clone: UCHL1

Isotype: Mouse IgG2a, κ


Workshop Number: IV N31

Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Common Marmoset

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE/Cy5 under optimal conditions. The solution is free of unconjugated PE/Cy5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with UCHL1 PE/Cy5

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: The UCHL1 antibody is commonly used in combination with antibodies against CD45RA to discern memory and naïve T cells. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁵ and formalin-fixed paraffin-embedded tissue sections⁴, Western blotting², and immunoprecipitation³.

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed for research use only.

- Application References:**
- Knapp W, et al. Eds. 1989. Leucocyte Typing IV. Oxford University Press. New York. (FC)
 - Ishii T, et al. 2001. *P. Natl. Acad. Sci. USA* 98:12138. (WB)
 - Ponsford M, et al. 2001. *Clin. Exp. Immunol.* 124:315. (IP)
 - Yamada M, et al. 1996. *Stroke* 27:1155. (IHC)
 - Sakkas LI, et al. 1998. *Clin. Diagn. Lab. Immunol.* 5:430. (IHC)
 - Baba N, et al. 2010. *Int. Immunol.* 22:237. PubMed
 - Thakral D, et al. 2008. *J. Immunol.* 180:7431. (FC) PubMed
 - Weiss L, et al. 2010. *P. Natl. Acad. Sci. USA* 107:10632. PubMed
 - Wu YY, et al. 2007. *Infect. Immun.* 75:4357. PubMed
 - Mozaffarian N, et al. 2008. *Rheumatology* 47:1335. PubMed
 - Roque S, et al. 2007. *J. Immunol.* 178:8028. PubMed
 - Yoshino N, et al. 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)

Description: CD45RO is a 180 kD single chain membrane glycoprotein. It is a splice variant of tyrosine phosphatase CD45, lacking the A, B, and C determinants. The CD45RO isoform is expressed on activated and memory T cells, some B cell subsets, activated monocytes/macrophages, and granulocytes. CD45RO enhances both T cell receptor and B cell receptor signaling mediated activation. CD45 and its isoforms non-covalently associate with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4. CD45 has also been reported to bind galectin-1 and CD22. CD45 isoform expression can change in response to cytokines.

- Antigen References:**
- Thomas M. 1989. *Annu. Rev. Immunol.* 7:339.
 - Trowbridge I, et al. 1994. *Annu. Rev. Immunol.* 12:85.

Related Products:

Product	Clone	Application
PE/Cy5 anti-human CD45RA	HI100	FC
PE/Cy5 anti-human CD45	HI30	FC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
PE anti-human CD45RB	MEM-55	FC
PE/Cy5.5 Mouse IgG2a, κ Isotype Ctrl	MOPC-173	FC, ICFC
Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.