

Product Data Sheet

Purified anti-human CD58

Catalog # / Size: 322502 / 100 µg

Clone: HCD58

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified HCD58, followed by anti-mouse IgGs FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Description: CD58, also known as lymphocyte function-associated antigen 3 (LFA-3) is a 45-70 kD cell surface protein that is a member of the immunoglobulin superfamily. Alternative splicing of CD58 gives rise to transmembrane and glycosylphosphatidylinositol (GPI)-anchored forms on cell surface. CD58 is expressed on both hematopoietic and non-hematopoietic cells including B cells, T cells, monocytes, erythrocytes, endothelial cells, epithelial cells, and fibroblasts. High levels are observed on memory T cells and dendritic cells. CD58 expressed on antigen presenting cells and target cells enhances T cell recognition via the binding of its cognate ligand, CD2, on the T cell surface. The HCD58 antibody recognizes human CD58 and has been shown to be useful for flow cytometry.

Antigen References:

1. Springer TA, *et al.* 1987. *Annu. Rev. Immunol.* 5:223.
2. Dustin ML, *et al.* 1987. *Nature* 329:846.
3. Arulanandam AR, *et al.* 1994. *J. Exp. Med.* 180:1861.
4. Sanders ME, *et al.* 1988. *J. Immunol.* 140:1401.

Related Products:

Product
 Purified Mouse IgG1, κ Isotype Ctrl
 APC Goat anti-mouse IgG (minimal x-reactivity)
 Biotin Goat anti-mouse IgG (minimal x-reactivity)
 PE Goat anti-mouse IgG (minimal x-reactivity)
 Cell Staining Buffer

Clone
 MOPC-21
 Poly4053
 Poly4053
 Poly4053

Application
 FC, ICFC, ICC, IF, IHC, IP, WB
 FC
 FC, ELISA, IHC, IF, WB
 FC
 FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.