


An Affymetrix Company

Anti-Mouse Granzyme B eFluor® 450

Catalog Number: 48-8898 Also known as: CGL1, CTLA-1

RUO: For Research Use Only. Not for use in diagnostic procedures.



Balb/c splenocytes were cultured with plate-bound Anti-Mouse CD3e Functional Grade Purified (cat. 16-0031) and Anti-Mouse CD28 Functional Grade Purified (cat. 16-0281) for 3 days, then cultured with Protein Transport Inhibitor Cocktail (cat. 00-4980) for an additional 5 hours. Cells were surface stained with Anti-Mouse CD8a PE (cat. 12-0081) followed by intracellular staining with 0.06 ug of Rat IgG2a K Isotype Control eFluor 450 (cat. 48-4321) (left) or 0.06 ug of Anti-Mouse Granzyme B eFluor 450 (right). Cells in the lymphocytes gate were used for analysis.

Product Information

Contents: Anti-Mouse Granzyme B eFluor®

450

Catalog Number: 48-8898

Clone: NGZB

Concentration: 0.2 mg/mL Host/Isotype: Rat IgG2a, kappa



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer **Temperature Limitation:** Store at 2-8°C. Do not

freeze. Light-sensitive material. **Batch Code:** Refer to vial

Use By: Refer to vial Contains Azide



Description

This NGZB monoclonal antibody reacts with mouse Granzyme B, which is a member of the granzyme serine protease family. Granzyme B is found in the granules of cytotoxic T cells and NK cells. Granzyme B has also been described as CGL1 (cathepsin G-like-1), a serine protease expressed only in cytotoxic T-lymphocytes after cell activation, and CTLA-1 (cytotoxic T lymphocyte-associated serine esterase 1) based on identification of mRNA in various cytotoxic T cells, but not observed in non-cytotoxic lymphoid cells. Granzyme B is crucial for the rapid induction of target cell death by apoptosis, induced by interaction with cytotoxic T cells. The receptor involved has been identified as mannose 6-phosphate receptor. This receptor functions as a death receptor for Granzyme B during cytotoxic T cell-induced apoptosis. This NGZB monoclonal antibody does not crossreact to human Granzyme B nor is staining blocked with GB11, suggesting it recognizes a different epitope.

Applications Reported

This NGZB antibody has been reported for use in intracellular staining and flow cytometric analysis.

Applications Tested

This NGZB antibody has been tested by intracellular staining followed by flow cytometric analysis of stimulated mouse splenocytes. This can be used at less than or equal to 0.125 μg per test. A test is defined as the amount (μg) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

eFluor® 450 is a replacement for Pacific Blue®. eFluor® 450 emits at 456 nm and is excited with the Violet laser (405 nm). Please make sure that your instrument is capable of detecting this fluorochome.

References

Kapp JA, Honjo K, Kapp LM, Xu XY, Cozier A, Bucy RP. TCR transgenic CD8+ T cells activated in the presence of TGFbeta express FoxP3 and mediate linked suppression of primary immune responses and cardiac allograft rejection. Int Immunol. 2006; 1-14.



Anti-Mouse Granzyme B eFluor® 450

Catalog Number: 48-8898 Also known as: CGL1, CTLA-1

RUO: For Research Use Only. Not for use in diagnostic procedures.

Smyth MJ, Trapani JA. Granzymes: exogenous proteinases that induce target cell apoptosis. Immunol Today. 1995 Apr;16(4):202-6.

Related Products

00-4980 Protein Transport Inhibitor Cocktail (500X)

00-8222 IC Fixation Buffer

00-8333 Permeabilization Buffer (10X)

12-0081 Anti-Mouse CD8a PE (53-6.7)

16-0031 Anti-Mouse CD3e Functional Grade Purified (145-2C11)

16-0281 Anti-Mouse CD28 Functional Grade Purified (37.51)

48-4321 Rat IgG2a K Isotype Control eFluor® 450 (eBR2a)