

Product Data Sheet

LEAF™ Purified anti-human CD85j (ILT2)

Catalog # / Size: 333704 / 500 µg

Clone: GHI/75

Isotype: Mouse IgG2b, κ

Workshop Number: V B032


Reactivity: Human

Preparation: The LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.

Formulation: 0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.1 EU/µg of the protein (<0.01 ng/µg of the protein) as determined by the LAL test.

Concentration: 1.0 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. This LEAF™ solution contains no preservative; handle under aseptic conditions.



Human peripheral blood lymphocytes stained with LEAF™ purified GHI/75, followed by anti-mouse IgG FITC

Applications:

Applications: FC - Quality tested
FA - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported application: Block HLA-G induced TGF-β1 production.

Application References: 1. Pulford K, *et al.* 1991. *Clin. Exp. Immunol.* 85:429
2. McIntire RH, *et al.* 2004. *J. Leukoc. Biol.* 76:1220

Description: CD85 is a group of Ig superfamily transmembrane glycoproteins called Ig-Like Transcripts (ILTs) or Leukocyte Immunoglobulin-like Receptors (LIRs). CD85j is the 110kD member, known as ILT2, LIR1, or LILRB1, and MIR7. ILT2 structurally has four Ig domains and contains ITIMs in its cytoplasmic tail that provide inhibitory signals by recruiting SHP-1. ILT2 is found on the surface of B cells, plasma cells, dendritic cells, monocytes, subsets of NK and T cells. The ligands of ILT2 include a broad range of HLA-A, -B molecules, some HLA-C and HLA-G molecules, and the human cytomegalovirus UL18 protein.

Antigen References: 1. Zola H, *et al.* 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers* Wiley-Liss A John Wiley & Sons Inc, Publication
2. Kirwan SE and Burshtyn DN. 2005. *J. Immunol.* 175:5006

Related Products: **Product**
LEAF™ Purified Mouse IgG2b, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
MG2b-57

Application
FC, ICFC, WB, IP, ICC, IF, FA
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.