

FITC anti-human CD71

Catalog # / Size: 334103 / 25 tests
334104 / 100 tests

Clone: CY1G4

Isotype: Mouse IgG2a, κ

Workshop Number: A015


Immunogen: NALM-6 pre-B cell line

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.5% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



PHA-stimulated (3 day) human peripheral blood lymphocytes stained with CY1G4 FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application References: 1. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
2. Park YS, *et al.* 2012. *Biochem Biophys Res Commun.* 423:19. PubMed.

Description: CD71 is a 95 kD type II homodimeric transmembrane glycoprotein also known as T9 and transferrin receptor. It is expressed on proliferating cells, reticulocytes, and erythroid precursors. CD71 plays a role in the control of cellular proliferation by facilitating the uptake of iron via ferrotransferrin binding and the recycling of apotransferrin to the cell surface.

Antigen References: 1. Hentze M, *et al.* 1996. *P. Natl. Acad. Sci. USA* 93:8175.
2. Trowbridge I, *et al.* 1993. *Annu. Rev. Cell Biol.* 9:129.

Related Products:

Product
FITC Rat IgG2a, κ Isotype Ctrl
Cell Staining Buffer

FITC Mouse IgG2a, κ Isotype Ctrl

Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone
RTK2758

MOPC-173

Application

FC, ICFC

FC, ICC, ICFC

FC, ICFC

FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.