

Product Data Sheet

PE/Cy7 anti-human CD95 (Fas)

Catalog # / Size: 305621 / 25 tests
305622 / 100 tests

Clone: DX2

Isotype: Mouse IgG1, κ

Workshop Number: VI C-64


Immunogen: CD95 transfected L cells

Reactivity: Human, **Cross-Reactivity:** African Green, Baboon, Capuchin Monkey, Chimpanzee, Common Marmoset, Cotton-topped Tamarin, Cynomolgus, Pigtailed Macaque, Rhesus, Sooty Mangabey

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with DX2 PE/Cy7

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: The DX2 antibody is useful for inducing apoptosis of Fas-positive cells. Additional reported applications (for the relevant formats) include: *in vitro* induction of apoptosis³ (DX2 antibody is required to be cross-linked for effective induction of apoptosis) and immunohistochemical staining^{4,5} of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded tissue sections. The LEAF™ Purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 305614).

Note: EOS9.1 antibody (cat. No. 305703/305704) can induce apoptosis without cross-linking process.

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed for research use only.

- Application References:**
- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. New York.
 - Cifone M, *et al.* 1994. *J. Exp. Med.* 180:1547. (Apop)
 - Zietz C, *et al.* 2001. *Am. J. Pathol.* 159:963. (IHC)
 - Sergi C, *et al.* 2000. *Am. J. Pathol.* 156:1589. (IHC)
 - Xie S, *et al.* 2010. *J. Immunol.* 184:2289. (FC) PubMed
 - Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 - Sestak K, *et al.* 2007. *Vet. Immunol. Immunopathol.* 119:21.
 - Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)

Description: CD95 is a 45 kD single chain type I glycoprotein also known as Fas, APO-1, and TNFRSF6. It is a member of the TNF receptor superfamily. CD95 is expressed on T and B lymphocytes, monocytes, neutrophils, and fibroblasts. CD95 expression is upregulated by activation. The extracellular region of CD95 binds to CD178 (Fas ligand). CD178 binding to CD95 induces apoptosis and has been shown to play a role in the maintenance of peripheral tolerance.

- Antigen References:**
- Krammer P, *et al.* 1994. *Immunol. Rev.* 142:175.
 - Nagata S, *et al.* 1995. *Science* 267:1449.

Related Products:

Product
 Cell Staining Buffer
 RBC Lysis Buffer (10X)
 PE/Cy7 Mouse IgG1, κ Isotype Ctrl
 Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone

MOPC-21

Application

FC, ICC, ICFC
 FC, ICFC
 FC, ICFC
 FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.