


Product Data Sheet

Purified anti-human CD150 (SLAM)

Catalog # / Size: 306302 / 100 µg
Clone: A12 (7D4)
Isotype: Mouse IgG1, κ
Immunogen: Activated human PBMC
Reactivity: Human, **Cross-Reactivity:** Baboon, Cynomolgus, Rhesus
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5 mg/ml
Storage: The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified A12, followed by biotinylated anti-mouse IgG and Sav-PE

Applications:

Applications: FC - *Quality tested*
 IHC, IP - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.5 µg per 10⁶ cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections, immunoprecipitation⁴, and costimulation^{1,5} of IFN-gamma production and T cell proliferation. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 306310).

Application References:

- Garcia V, *et al.* 2001. *J. Immunol.* 167:5719. (Costim)
- Vincent S, *et al.* 2002. *J. Virol.* 76:6121.
- Cocks B, *et al.* 1995. *Nature* 376:260.
- Sayos J, *et al.* 2001. *Blood* 97:3867. (IP)
- Aversa G, *et al.* 1997. *J. Immunol.* 158:4036. (Costim)
- Spencer M, *et al.* 2010. *Am. J. Physiol Endocrinol Metab.* 299:1016. PubMed
- Baldanzi G., *et al.* 2011. *J. Immunol.* 187:5941. PubMed.

Description: CD150 is a 70-95 kD type I transmembrane glycoprotein also known as SLAM or IPO-3. It is a member of the Ig superfamily. It is expressed on a subset of T cells, B cells, dendritic cells, and endothelial cells. The expression of CD150 is upregulated upon activation. CD150 binds to itself as the ligand to be involved in B cell costimulation, proliferation, immunoglobulin production, and signal transduction.

Antigen References:

- Cocks B, *et al.* 1995. *Nature* 376:260.
- Pinchouk V, *et al.* 1988. *AntiCancer Res.* 8:1377.
- Polacino P, *et al.* 1996. *J. Med. Primatol.* 25:201.
- Punnonen J, *et al.* 1997. *J. Exp. Med.* 185:993.

Related Products:

Product
 Purified Mouse IgG1, κ Isotype Ctrl
 APC Goat anti-mouse IgG (minimal x-reactivity)
 Biotin Goat anti-mouse IgG (minimal x-reactivity)
 PE Goat anti-mouse IgG (minimal x-reactivity)
 PE Streptavidin
 Cell Staining Buffer
 RBC Lysis Buffer (10X)

Clone
 MOPC-21
 Poly4053
 Poly4053
 Poly4053

Application
 FC, ICFC, ICC, IF, IHC, IP, WB
 FC
 FC, ELISA, IHC, IF, WB
 FC
 FC, ICFC
 FC, ICC, ICFC
 FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.