

Purified anti-human CD134 (OX40)

Catalog # / Size: 350001 / 25 µg
350002 / 100 µg

Clone: Ber-ACT35 (ACT35)

Isotype: Mouse IgG1, κ

Immunogen: HTLV 1-transformed HUT 102 cells


Reactivity: Human, **Cross-Reactivity:** Chimpanzee

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes were activated for 3 days with PHA, then stained with OX40 (clone Ber-ACT35) PE (filled histogram) or mouse IgG1, κ PE isotype control (open histogram).

Applications:

Applications: FC - *Quality tested*
ELISA, IHC, WB - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting¹, immunoprecipitation¹, immunohistochemical staining^{2,3} of paraffin embedded⁷ and frozen tissue sections, ELISA⁴, and functional assay⁵. The LEAF™ or Ultra-LEAF™ purified antibody is recommended for functional assays (contact our custom solutions team).

Application References:

1. Latza U, *et al.* 1994. *Eur. J. Immunol.* 24:677. (WB, IP)
2. Durkop H, *et al.* 1995. *Brit. J. Haematol.* 91:927. (IHC)
3. Durkop H, *et al.* 1997. *Brit. J. Haematol.* 98:863. (IHC)
4. Willett B, *et al.* 2007. *J. Virol.* 81:9665. (ELISA)
5. Li M and Zhang Y. *et al.* 2005. *Cell. Mol. Immunol.* 2:467. (FA)
6. Glociczki ML, *et al.* 2012. *Clin J Am Soc Nephrol.* PubMed
7. Domingos PL, *et al.* 2012. *An. Bras. Dermatol.* 87:851. (IHC)

Description: CD134, also known as OX40 and TNFRSF14, is a 50 kD type I transmembrane glycoprotein. It is a member of the TNF receptor family. OX40 is expressed on activated T lymphocytes including Th1, Th2, Th17, and Treg cells. The interaction of OX40 with OX40L results in B cell proliferation and antibody secretion, regulation of primary T cell expansion, and T cell survival. OX40 influences the size of the T cell memory pool and regulation of CD4⁺ T cell tolerance.

Antigen References:

1. Smith CA, *et al.* 1994. *Cell* 76:959.
2. Chen AL, *et al.* 1999. *Immunity* 11:689.
3. Croft M. 2010. *Annu. Rev. Immunol.* 28:57.
4. Ruby CE, *et al.* 2009. *J. Immunol.* 183:5079.
5. Klinger M, *et al.* 2009. *J. Immunol.* 182:4581.

Related Products:

Product
Purified Mouse IgG1, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
MOPC-21

Application
FC, ICFC, ICC, IF, IHC, IP, WB
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.